

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human LRP-6 in direct ELISAs. In direct ELISAs, approximately 5% cross-reactivity with recombinant mouse (rm) LRP-6 and no cross-reactivity with recombinant human (rh) LRP-1, -4, -5, rhLRP-6 (intracellular domain), or rhLDL R is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 255302
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human LRP-6 Ala20-Pro1368 Accession # O75581
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
Immunocytochemistry	8-25 µg/mL	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA

<p>Flow Cytometry</p> <p>Detection of LRP-6 in MDA-MB-231 Human Cell Line by Flow Cytometry. MDA-MB-231 human breast cancer cell line was stained with Mouse Anti-Human LRP-6 Monoclonal Antibody (Catalog # MAB1505, filled histogram) or isotype control antibody (Catalog # MAB003, open histogram), followed by Phycoerythrin-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # F0102B).</p>	<p>Immunocytochemistry</p> <p>LRP-6 in DU145 Human Cell Line. LRP-6 was detected in immersion fixed DU145 human prostate carcinoma cell line using Mouse Anti-Human LRP-6 Monoclonal Antibody (Catalog # MAB1505) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). Specific staining was localized to cell surfaces and cytoplasm. View our protocol for Fluorescent ICC Staining of Cells on Coverslips.</p>
--	---

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

LRP-6 (Low-density lipoprotein receptor-related protein 6) is a 200-210 kDa member of the LDLR family of proteins. It is widely expressed, and serves as a coreceptor for both Wnt and parathyroid hormone. In the Wnt system, LRP-6 associates with select Fzd multipass receptors; in the PTH system, LRP-6 complexes with PTH1R. Mature human LRP-6 is a 1594 amino acid (aa) type I transmembrane glycoprotein. It contains a 1351 aa extracellular region (aa 20-1370) plus a 220 aa cytoplasmic domain (aa 1394-1613). The cytoplasmic domain contains two palmitoylation sites, one ubiquitination residue, and multiple phosphorylation motifs.