

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human MICB in direct ELISAs and Western blots. Does not cross-react with recombinant human MICA.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 236511
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human MICB Ala23-Gly298 Accession # CA118747
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	Recombinant Human MICB Fc Chimera, aa 23-298 (Catalog # 1599-MB)
<b>Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below
<b>Human MICB Sandwich Immunoassay</b>		<b>Reagent</b>
<b>ELISA Capture</b>	2-8 µg/mL	Human MICB Antibody (Catalog # MAB1599)
<b>ELISA Detection</b>	0.1-0.4 µg/mL	Human MICB Biotinylated Antibody (Catalog # BAF1599)
<b>Standard</b>		Recombinant Human MICB Fc Chimera (Catalog # 1599-MB)
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

MICB (MHC class I chain-related gene B) is a transmembrane glycoprotein that functions as a ligand for NKG2D. A closely related protein, MICA, shares 85% amino acid identity with MICB. These 2 proteins are distantly related to the MHC class I proteins. MICA and MICB (MICA/B) possess three extracellular immunoglobulin-like domains, but have no capacity to bind peptide or interact with  $\beta$ 2-microglobulin. The genes encoding MICA/B are found within the major histocompatibility complex on human chromosome 6. The MICB locus is polymorphic with more than 15 recognized human alleles. MICA/B are minimally expressed on normal cells, but are frequently expressed on epithelial tumors and can be induced by bacterial and viral infections. MICA/B are ligands for NKG2D, an activating receptor expressed on NK cells, NKT cells,  $\gamma\delta$  T cells, and CD8<sup>+</sup>  $\alpha\beta$  T cells. Recognition of MICA/B by NKG2D results in the activation of cytolytic activity and/or cytokine production by these effector cells. MICA/B recognition is involved in tumor surveillance, viral infections, and autoimmune diseases. The release of soluble forms of MICA/B from tumors down-regulates NKG2D surface expression on effector cells resulting in the impairment of anti-tumor immune response (1-7).

**References:**

1. Groh, V. *et al.* (2001) *Nature Immunol.* **2**:255.
2. Stephens, H. (2001) *Trends Immunol.* **22**:378.
3. Bauer, S. *et al.* (1999) *Science* **285**:727.
4. Groh, V. *et al.* (2002) *Nature* **419**:734.
5. Steinle, A. *et al.* (2001) *Immunogenetics* **53**:279.
6. Pende, D. *et al.* (2002) *Cancer Res.* **62**:6178.
7. Salih, H. *et al.* (2003) *Blood* **102**:1389.