



DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human CXCR4.
Source	Recombinant Monoclonal Mouse IgG _{2A} Clone # 12G5R
Purification	Protein A or G purified from cell culture supernatant
Immunogen	CP-MAC-infected SUP-T1 human T cell lymphoblastic lymphoma line Accession # P61073
Formulation	Supplied as a solution in PBS. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS	
Please Note: Optimal dilutions should be determined by each laboratory for each application. <i>General Protocols</i> are available in the <i>Technical Information</i> section on our website.	
	Recommended Concentration Sample
Flow Cytometry	0.25 µg/10 ⁶ cells See Below
Neutralization	Measured by its ability to neutralize CXCL12/SDF-1α-induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CXCR4. The Neutralization Dose (ND ₅₀) is typically 0.3-1.2 µg/mL in the presence of 1 ng/mL Recombinant Human/Feline/Rhesus Macaque CXCL12/SDF-1α.

DATA	
<p>Flow Cytometry</p>  <p>Detection of CXCR4 in Human Blood Lymphocytes by Flow Cytometry. Human peripheral blood lymphocytes were stained with Mouse Anti-Human CD19 APC-conjugated Monoclonal Antibody (Catalog # FAB4867A) and either (A) Mouse Anti-Human CXCR4 Monoclonal Antibody (Catalog # MAB170R) or (B) Mouse IgG_{2A} Isotype Control (Catalog # MAB003) followed by Phycoerythrin-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # F0102B). View our protocol for Staining Membrane-associated Proteins.</p>	<p>Flow Cytometry</p>  <p>Chemotaxis Induced by CXCL12/SDF-1α and Neutralization by Human CXCR4 Antibody. Recombinant Human/Feline/Rhesus Macaque CXCL12/SDF-1α (Catalog # 350-NS) chemoattracts the BaF3 mouse pro-B cell line transfected with human CXCR4 in a dose-dependent manner (orange line). The amount of cells that migrated through to the lower chemotaxis chamber was measured by Resazurin (Catalog # AR002). Chemotaxis elicited by Recombinant Human/Feline/Rhesus Macaque CXCL12/SDF-1α (1 ng/mL) is neutralized (green line) by increasing concentrations of Mouse Anti-Human CXCR4 Monoclonal Antibody (Catalog # MAB170R). The ND₅₀ is typically 0.3-1.2 µg/mL.</p>

PREPARATION AND STORAGE	
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C, as supplied. • 1 month, 2 to 8 °C under sterile conditions after opening. • 6 months, -20 to -70 °C under sterile conditions after opening.

BACKGROUND	
CXCR4 is a G-protein-linked seven transmembrane spanning receptor that binds stromal cell-derived factor-1 (SDF-1). CXCR4 acts as a co-factor for T-cell tropic HIV-1 and -2 viral entry into cells.	
References:	
1. Endres, M.J. <i>et al.</i> (1996) <i>Cell</i> 87 :745.	