

Mouse Follistatin-like 1/FSTL1 Antibody

Recombinant Mouse Follistatin-like 1/FSTL1 under non-reducing conditions only

Monoclonal Rat IgG_{2A} Clone # 228210 Catalog Number: MAB1738

DESCRIPTION	
Species Reactivity	Mouse
Specificity	Detects mouse Follistatin-like 1/FSTL1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, 100% cross-reactivity with recombinant human Follistatin-like 1/FSTL1 is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 228210
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Follistatin-like 1/FSTL1 Glu19-Ile306 Accession # Q62356
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.
APPLICATIONS	
Please Note: Optimal diluti	ons should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.
	Recommended Sample Concentration

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 µg/mL

BACKGROUND

Western Blot

Follistatin-Like 1, also known as FSTL1 and follistatin-related protein (FRP), is a secreted protein that contains a follistatin-like domain, a Kazal-like domain and a von Willebrand factor C domain. Human FSTL1 shares 92% amino acid sequence identity with the mouse FSTL1, also known as TSC-36.

biotechne