

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human DCIR in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 216110
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human DCIR isoform 1 Gln70-Leu237 Accession # Q9UMR7
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Human DCIR/CLEC4A under non-reducing conditions only
Flow Cytometry	2.5 µg/10 ⁶ cells	Human peripheral blood neutrophils
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

DCIR, also known as Lectin-like Immunoreceptor (LLIR), is a type II membrane protein belonging to the C-type lectin domain family and is designated CLEC4A (previously designated CLECSF6). Four transcript variants encoding distinct isoforms have been identified. DCIR contains one carbohydrate recognition domain in its C-terminal extracellular domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM) in its cytoplasmic domain. Besides DC, DCIR is expressed on B cells, monocytes/macrophages and granulocytes.