

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TIM-1/KIM-1/HAVCR in direct ELISAs and Western blots. Does not cross-react with recombinant mouse (rm) TIM-1, rmTIM-2, or rhTIM-3.
Source	Monoclonal Mouse IgG _{2B} Clone # 219211
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human TIM-1/KIM-1/HAVCR Ser21-Thr288 Accession # AAC39862
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
Immunohistochemistry	8-25 µg/mL	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	


DATA

Western Blot




Detection of Human TIM-1/KIM-1/HAVCR by Western Blot. Western blot shows lysates of human CD4⁺ cells treated (+) with 5 µg/mL of Hamster Anti-Mouse CD3ε Monoclonal Antibody (Catalog # [MAB484](#)) and 1 µg/mL of Rat Anti-Mouse CD28 Monoclonal Antibody (Catalog # [MAB4831](#)) for 24 hours. PVDF membrane was probed with 1 µg/mL of Mouse Anti-Human TIM-1/KIM-1/HAVCR Monoclonal Antibody (Catalog # [MAB1750](#)) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # [HAF007](#)). A specific band was detected for TIM-1/KIM-1/HAVCR at approximately 80 kDa (as indicated). This experiment was conducted under reducing conditions and using [Immunoblot Buffer Group 1](#).

Flow Cytometry



Detection of TIM-1/KIM-1/HAVCR in Th2-stimulated Human PBMCs by Flow Cytometry. (A) Unstimulated and (B) Th2-stimulated human PBMCs were stained with Mouse Anti-Human TIM-1/KIM-1/HAVCR Monoclonal Antibody (Catalog # [MAB1750](#)) followed by Allophycocyanin-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # [F0101B](#)) and Human CD4 PerCP-conjugated Monoclonal Antibody (Catalog # [FAB3791C](#)). Quadrant markers were set based on control antibody staining (Catalog # [MAB0041](#)).

Immunohistochemistry



TIM-1/KIM-1/HAVCR in Human Kidney. TIM-1/KIM-1/HAVCR was detected in immersion fixed paraffin-embedded sections of human kidney using 25 µg/mL Mouse Anti-Human TIM-1/KIM-1/HAVCR Monoclonal Antibody (Catalog # [MAB1750](#)) overnight at 4 °C. Tissue was stained with the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # [CTS002](#)) and counterstained with hematoxylin (blue). View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

TIM-1 (T cell-immunoglobulin-mucin; also KIM-1 and HAVcr-1) is a 100 kDa, type I transmembrane glycoprotein member of the TIM family of immunoglobulin superfamily molecules (1-3). This gene family is involved in the regulation of Th1 and Th2-cell-mediated immunity. Human TIM-1 is synthesized as a 359 amino acid (aa) precursor that contains a 20 aa signal sequence, a 270 aa extracellular domain (ECD), a 21 aa transmembrane segment and a 48 aa cytoplasmic domain (4-6). The ECD contains one V-type Ig-like domain and a mucin region characterized by multiple PTTTTL motifs. The mucin region undergoes extensive O-linked glycosylation. The TIM-1 gene is highly polymorphic and undergoes alternate splicing (1). For instance, the presence of a six aa sequence (MTTTPV) at position #137 of the mature molecule is associated with protection from atopy in people with a history of hepatitis A (7, 8). There are two cytoplasmic alternate splice forms of TIM-1. One is a long (359 aa) kidney form termed TIM-1b, and one is a short (334 aa) liver form termed TIM-1a. Both are identical through the first 323 aa of their precursors. TIM-1b contains a tyrosine phosphorylation motif that is not present in 1a (6). TIM-1 is also known to circulate as a soluble form. Constitutive cleavage by an undefined MMP (possibly ADAM33) releases an 85 - 90 kDa soluble molecule (6). The ECD of human TIM-1 is 50% and 43% aa identical to mouse and canine TIM-1 ECD, respectively. The only two reported ligands for TIM-1 are TIM-4 and the hepatitis A virus (4, 9). However, others are believed to exist, and based on the ligand for TIM-3, one may well be an S-type lectin (10). TIM-1 ligation induces T cell proliferation and promotes cytokine production (1, 10).

References:

1. Meyers, J.H. *et al.* (2005) Trends Mol. Med. **11**:1471.
2. Kuchroo, V.K. *et al.* (2003) Nat. Rev. Immunol. **3**:454.
3. Mariat, C. *et al.* (2005) Phil. Trans. R. Soc. B **360**:1681.
4. Feigelstock, D. *et al.* (1998) J. Virol. **72**:6621.
5. Ichimura, T. *et al.* (1998) J. Biol. Chem. **273**:4135.
6. Bailly, V. *et al.* (2002) J. Biol. Chem. **277**:39739.
7. Umetsu, D.T. *et al.* (2005) J. Pediatr. Gastroenterol. Nutr. **40**:S43.
8. Gao, P-S. *et al.* (2005) J. Allergy Clin. Immunol. **115**:982.
9. Zhu, C. *et al.* (2005) Nat. Immunol. **6**:1245.
10. Meyers, J.H. *et al.* (2005) Nat. Immunol. **6**:455.

PRODUCT SPECIFIC NOTICES

This product is covered by one or more of the following US Patents 7,300,652; 7,041,290; 6,664,385 and other US and foreign patents pending or issued.