

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human p53 that is phosphorylated at S15 in Western blots. In Western blots, no cross-reactivity with human p53 that is unphosphorylated at S15 is detected.
Source	Monoclonal Mouse IgG ₁ Clone # 261352
Purification	Protein A or G from hybridoma culture supernatant
Immunogen	Phosphopeptide containing the human p53 S15 site Accession # P04637
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1-2 µg/mL	See Below
Intracellular Staining by Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CytoF-reported	Brodie, T.M. <i>et al.</i> (2018) <i>Cytometry Part A</i> . 93 : 406. Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

The p53 tumor suppressor protein acts to enforce cell cycle checkpoints or signal apoptosis in cells that have incurred genotoxic damage. The ATM or ATR kinases can phosphorylate p53 at serine 15 (S15), which leads to cell cycle arrest. Serine 15 phosphorylation leads to p53 stabilization and enhances transactivation of p53 target genes.