

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human BMP-1/PCP in direct ELISAs and Western blots.
Source	Monoclonal Rat IgG _{2B} Clone # 264822
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human BMP-1/PCP Ala121-Gln730 Accession # NP_001190
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	2 µg/mL	See Below
Immunoprecipitation	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human BMP-1/PCP (Catalog # 1927-ZN), see our available Western blot detection antibodies

DATA

Western Blot



Detection of Human BMP-1/PCP by Western Blot. Western blot shows lysates of Jurkat human acute T cell leukemia cell line and human skin tissue. PVDF Membrane was probed with 2 µg/mL of Human BMP-1/PCP Monoclonal Antibody (Catalog # MAB1927) followed by HRP-conjugated Anti-Rat IgG Secondary Antibody (Catalog # HAF005). A specific band was detected for BMP-1/PCP at approximately 180 kDa (as indicated). This experiment was conducted under non-reducing conditions and using [Immunoblot Buffer Group 1](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Bone morphogenetic protein 1 (BMP-1), also known as procollagen C-proteinase (PCP), is a zinc protease of the astacin family (1, 2). BMP-1/PCP plays a key role in formation of extracellular matrix (ECM) by converting precursor proteins into their mature and functional forms. The precursor proteins identified as substrates for BMP-1/PCP include collagens, biglycan, laminin 5, dentin matrix protein-1, and lysyl oxidase (3). There are six alternatively spliced forms known to be derived from the BMP-1 gene, and isoform 1 consisting of residues 1 to 730 was expressed. The secreted and purified protein does not contain the signal peptide (amino acid residues 1-22) and pro domain (residues 23-120), but contain protease (residues 121-321), CUB I (residues 322-434), CUB II (residues 435-546), EGF-like (residues 547-588) and CUB III (residues 591-703) domains. The pro domain is apparently cleaved by a furin-like proprotein convertase (4). The purified BMP-1/PCP is an active protease and its peptidase activity can be determined as described above. The purified BMP-1/PCP is predicted to possess procollagen C-proteinase activity because it contains the minimal domain structure required (5).

References:

1. Wozney, J.M. et al. (1988) Science **242**:1528.
2. Bond, J.S. and R.J. Beynon (1995) Protein Sci. **4**:1247.
3. Steiglitz, B.M. et al. (2004) J. Biol. Chem. **279**:980.
4. Leighton, M. and K.E. Kadler (2003) J. Biol. Chem. **278**:18478.
5. Hartigan, N. et al. (2003) J. Biol. Chem. **278**:18045.