

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human MRP1. Detects human MRP1-transfected cells, but not irrelevant transfectants by flow cytometry.
<b>Source</b>	Monoclonal Mouse IgG <sub>2A</sub> Clone # 464510
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	HEK293 human embryonic kidney cell line transfected with human MPR1 Met1-Val1531 Accession # P33527
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Intracellular Staining by Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

MRP1 (Multidrug Resistance associated Protein 1), also known as ABCC1, belongs to the family of ATP-Binding Cassette (ABC) transporters. It confers multidrug resistance on tumor cells and is overexpressed in various human multidrug resistant tumor cell lines.