

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human GDF-11/BMP-11 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 937320
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human GDF-11/BMP-11 Asn299-Ser407 Accession # O95390
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below

DATA

Immunocytochemistry



GDF-11/BMP-11 in HEK293 Human Cell Line. GDF-11/BMP-11 was detected in immersion fixed HEK293 human embryonic kidney cell line transfected with GDF-11 using Mouse Anti-Human GDF-11/BMP-11 Monoclonal Antibody (Catalog # MAB19583) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Growth Differentiation Factor 11 (GDF-11), also known as BMP-11, is a member of the TGF- β superfamily and is highly related to GDF-8. GDF-11 encodes a 407 amino acid (aa) prepropeptide which contains a signal sequence for secretion and an RXXR proteolytic processing site to yield a 109 aa residue carboxy-terminal mature protein (1). Mature GDF-11 contains the canonical 7-cysteine motif common to other TGF- β superfamily members; however, like the TGF- β s, Activins and GDF-8, GDF-11 also contains one extra pair of cysteine residues. At the amino acid sequence level, mature human, mouse, rat and chicken GDF-11 are 99-100% identical. GDF-11 and GDF-8 share 90% amino acid sequence identity within the mature protein. As detected by in situ hybridization, GDF-11 is expressed in diverse regions of the mouse embryo: tailbud, somitic precursors, limbs, mandibular and branchial arches, dorsal neural tube, odontoblasts, nasal epithelium, and particular regions of the brain (1, 2). Targeted deletion of GDF-11, in mice, results in a spectrum of abnormalities including palatal malformation, vertebral defects, elongated trunks with a reduced or absent tail, missing or malformed kidneys, and an increased number of neurons in the olfactory epithelium (2-5). GDF-11 signals through the Activin type II receptors and induces phosphorylation of Smad2 to mediate axial patterning (6). Systemic GDF-11 levels decline with age and administration of higher levels of GDF-11 can reverse age-related cardiac hypertrophy (7). In addition, systemic administration of recombinant GDF-11 protein restores genomic integrity and health of muscle stem cells, neurovasculature and enhances neurogenesis (8, 9).

References:

1. Gamer, L.W. *et al.* (1999) *Dev. Biol.* **208**: 222.
2. Nakashima, M. *et al.* (1999) *Mech. Dev.* **80**:185.
3. Gad, J.M. and P.P.L. Tam (1999) *Curr. Biol.* **9**:R783.
4. McPherron, A.C. *et al.* (1999) *Nat. Genet.* **22**:260.
5. Esquela, A.F. and S.J. Lee (2003) *Dev. Biol.* **257**:356.
6. Oh, S.P. *et al.* (2002) *Genes & Dev.* **16**:274.
7. Loffredo, F.S. *et al.* (2013) *Cell.* **153**:828.
8. Katsimpardi, L. *et al.* (2014) *Science* (ahead of print).
9. Sinha, M. *et al.* (2014) *Science* (ahead of print).