




| DESCRIPTION               |   |
|---------------------------|---|
| <b>Species Reactivity</b> | Human   |
| <b>Specificity</b>        | Detects human CCR7.   |
| <b>Source</b>             | Monoclonal Mouse IgG <sub>2A</sub> Clone # 150503   |
| <b>Purification</b>       | Protein A or G purified from hybridoma culture supernatant  |
| <b>Immunogen</b>          | Human CCR7 transfectants<br>Met1-Pro378<br>Accession # AAA58615   |
| <b>Endotoxin Level</b>    | <0.10 EU per 1 µg of the antibody by the LAL method.  |
| <b>Formulation</b>        | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.<br>*Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS. |

| APPLICATIONS   |   |               |
|--|---|---------------|
| <b>Please Note:</b> Optimal dilutions should be determined by each laboratory for each application. <i>General Protocols</i> are available in the <i>Technical Information</i> section on our website. |   |               |
|  | <b>Recommended Concentration</b>  | <b>Sample</b> |
| <b>Flow Cytometry</b>  | 0.25 µg/10 <sup>6</sup> cells   | See Below     |
| <b>Immunocytochemistry</b>   | 8-25 µg/mL  | See Below     |
| <b>CyTOF-reported</b>  | Mei, H.E. <i>et al.</i> (2015) <i>J. Immunol.</i> <b>194</b> : 2022. Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.   |               |
| <b>Neutralization</b>  | Measured by its ability to neutralize CCL19/MIP-3β-induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CCR7. The Neutralization Dose (ND <sub>50</sub> ) is typically 1-5 µg/mL in the presence of 50 ng/mL Recombinant Human CCL19/MIP-3β. |               |

| DATA  |   |
|---|---|
| <p><b>Immunocytochemistry</b></p>  <p><b>CCR7 in Human PBMCs.</b><br/>CCR7 was detected in immersion fixed human peripheral blood mononuclear cells (PBMCs) using 25 µg/mL Mouse Anti-Human CCR7 Monoclonal Antibody (Catalog # MAB197) for 3 hours at room temperature. Cells were stained (red) and counterstained (green). View our protocol for <a href="#">Fluorescent ICC Staining of Non-adherent Cells</a>.</p> | <p><b>Flow Cytometry</b></p>  <p><b>Detection of CCR7 in Human PBMCs by Flow Cytometry.</b><br/>Human peripheral blood mononuclear cells (PBMCs) were stained with Mouse Anti-Human CD4 PE-conjugated Monoclonal Antibody (Catalog # FAB3791P) and either (A) Mouse Anti-Human CCR7 Monoclonal Antibody (Catalog # MAB197) or (B) Mouse IgG<sub>2A</sub> Isotype Control (Catalog # MAB003) followed by Allophycocyanin-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # F0101B). View our protocol for <a href="#">Staining Membrane-associated Proteins</a>.</p> |



## PREPARATION AND STORAGE

**Reconstitution** Reconstitute at 0.5 mg/mL in sterile PBS.

**Shipping** The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.  
\*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C

**Stability & Storage** Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 6 months, -20 to -70 °C under sterile conditions after reconstitution.

## BACKGROUND

CCR7 is a 7 transmembrane G protein coupled chemokine receptor. CCR7 is expressed on T cells and mature dendritic cells and transduces chemotactic signals in response to CCL19 and CCL21. Human CCR7 shares 87% amino acid sequence identity with mouse CCR7.