

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human IL-1 $\alpha$ /IL-1F1 in ELISAs and Western blots. In ELISAs, this antibody does not cross-react with recombinant human (rh) IL-1 $\beta$ , -2, -3, -4, -5, -6, -7, -8, -9, -10, -11, -12, -13, rIL-1 $\alpha$ , -1 $\beta$ , -3, -4, -5, -6, -7, -9, or -13.
<b>Source</b>	Monoclonal Mouse IgG <sub>2A</sub> Clone # 4414
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human IL-1 $\alpha$ /IL-1F1 Ser113-Ala271 Accession # P01583
<b>Endotoxin Level</b>	<0.10 EU per 1 $\mu$ g of the antibody by the LAL method.
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 $\mu$ m filtered solution in PBS.


## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the Technical Information section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 $\mu$ g/mL	Recombinant Human IL-1 $\alpha$ /IL-1F1 (Catalog # 200-LA)
<b>Immunocytochemistry</b>	8-25 $\mu$ g/mL	See Below
<b>Human IL-1<math>\alpha</math>/IL-1F1 Sandwich Immunoassay</b>		<b>Reagent</b>
<b>ELISA Capture</b>	2-8 $\mu$ g/mL	Human IL-1 $\alpha$ /IL-1F1 Antibody (Catalog # MAB200)
<b>ELISA Detection Standard</b>	0.1-0.4 $\mu$ g/mL	Human IL-1 $\alpha$ /IL-1F1 Biotinylated Antibody (Catalog # BAF200) Recombinant Human IL-1 $\alpha$ /IL-1F1 (Catalog # 200-LA)
<b>Neutralization</b>	Measured by its ability to neutralize IL-1 $\alpha$ /IL-1F1-induced proliferation in the D10.G4.1 mouse helper T cell line. Symons, J.A. <i>et al.</i> (1987) in Lymphokines and Interferons, a Practical Approach. Clemens, M.J. <i>et al.</i> (eds): IRL Press. 272. The Neutralization Dose (ND <sub>50</sub> ) is typically 0.05-0.15 $\mu$ g/mL in the presence of 0.05 ng/mL Recombinant Human IL-1 $\alpha$ /IL-1F1.	


## DATA

### Immunocytochemistry



**IL-1 $\alpha$ /IL-1F1 in Human PBMCs.** IL-1 $\alpha$ /IL-1F1 was detected in immersion fixed human peripheral blood mononuclear cells (PBMCs) using 10  $\mu$ g/mL Human IL-1 $\alpha$ /IL-1F1 Monoclonal Antibody (Catalog # MAB200) for 3 hours at room temperature. Cells were stained with the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). View our protocol for Fluorescent ICC Staining of Non-adherent Cells.

### Neutralization



**Cell Proliferation Induced by IL-1 $\alpha$ /IL-1F1 and Neutralization by Human IL-1 $\alpha$ /IL-1F1 Antibody.** Recombinant Human IL-1 $\alpha$ /IL-1F1 (Catalog # 200-LA) stimulates proliferation in the D10.G4.1 mouse helper T cell line in a dose-dependent manner (orange line). Proliferation elicited by Recombinant Human IL-1 $\alpha$ /IL-1F1 (0.05 ng/mL) is neutralized (green line) by increasing concentrations of Human IL-1 $\alpha$ /IL-1F1 Monoclonal Antibody (Catalog # MAB200). The ND<sub>50</sub> is typically 0.05-0.15  $\mu$ g/mL.

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Interleukin 1 (IL-1) is a name that designates two proteins, IL-1 $\alpha$  and IL-1 $\beta$ , which are the products of distinct genes, but which show approximately 25% amino acid sequence identity and which recognize the same cell surface receptors. Although IL-1 production is generally considered to be a consequence of inflammation, recent evidence suggests that IL-1 is also temporarily upregulated during bone formation and the menstrual cycle and can be induced in response to nervous system stimulation. In response to classic stimuli produced by inflammatory agents, infections or microbial endotoxins, a dramatic increase in the production of IL-1 by macrophages and various other cells is seen. Cells in particular known to produce IL-1 include osteoblasts, monocytes, macrophages, keratinocytes, Kupffer cells, hepatocytes, thymic and salivary gland epithelium, Schwann cells, fibroblasts, and glia (oligodendroglia, astrocytes, and microglia).

IL-1 $\alpha$  and IL-1 $\beta$  are both synthesized as 31 kDa precursors that are subsequently cleaved into proteins with molecular weights of approximately 17,000 Da. Neither precursor contains a typical hydrophobic signal peptide sequence and most of the precursor form of IL-1 $\alpha$  remains in the cytosol of cells, although there is evidence for a membrane-bound form of the precursor form of IL-1 $\alpha$ . The IL-1 $\alpha$  precursor reportedly shows full biological activity in the EL-4 assay. Among various species, the amino acid sequence of mature IL-1 $\alpha$  is conserved 60% to 70% and human IL-1 has been found to be biologically active on murine cell lines. Both forms of IL-1 bind to the same receptors, designated type I and type II. Evidence suggests that only the type I receptor is capable of signal transduction and that the type II receptor may function as a decoy, binding IL-1 and thus preventing binding of IL-1 to the type I receptor.