

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human DEC-205/CD205 in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant mouse DEC-205 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 523218
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human DEC-205/CD205 Cys216-Cys501 Accession # O60449
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	2 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

DEC-205, also known as CD205 and lymphocyte antigen 75 (Ly 75), is a type I transmembrane protein that is primarily expressed on dendritic cells and thymic epithelial cells. The extracellular region of DEC-205 contains ten C-type lectin-like domains, a fibronectin type II domain and a ricin B-type lectin domain. DEC-205 functions as an endocytic receptor for antigens. The recombinant protein used to generate the anti-human DEC-205 antibody contains the first two C-type lectin domains.