


DESCRIPTION	
<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse Complement Component C5a in Western blots. In Western blots, this antibody does not cross-react with recombinant mouse Complement Component C3d, recombinant human (rh) Complement Component C3a, or rhComplement Component C5a.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 295108
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant mouse Complement Component C5a Asn679-Arg755 Accession # P06684
<b>Endotoxin Level</b>	<0.10 EU per 1 µg of the antibody by the LAL method.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 µg/mL	Recombinant Mouse Complement Component C5a (Catalog # 2150-C5)
<b>Mouse Complement Component C5a Sandwich Immunoassay</b>		<b>Reagent</b>
<b>ELISA Capture</b>	2-8 µg/mL	Mouse Complement Component C5a Antibody (Catalog # MAB21501)
<b>ELISA Detection Standard</b>	0.1-0.4 µg/mL	Mouse Complement Component C5a Biotinylated Antibody (Catalog # BAF2150) Recombinant Mouse Complement Component C5a (Catalog # 2150-C5)
<b>Neutralization</b>		Measured by its ability to neutralize Complement Component C5a-induced N-acetyl-β-D-glucosaminidase release in the dibutyryl cyclic-AMP differentiated U937 human histiocytic lymphoma cell line. Klos, A. <i>et al.</i> (1992) <i>Biochemistry</i> 31:11274. The Neutralization Dose (ND <sub>50</sub> ) is typically 0.1-0.5 µg/mL in the presence of 50 ng/mL Recombinant Mouse Complement Component C5a and cytochalasin-B.

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Mouse Complement 5a (C5a) is an enzymatically generated glycoprotein that belongs to a family of structurally and functionally related proteins known as Anaphylatoxins. C5a is a 77 amino acid (aa) peptide that is created by the activity of C5a convertase on the C5  $\alpha$ -chain (1, 2). Mouse C5a has four  $\alpha$ -helices, plus three intrachain disulfide bonds that create a triple loop structure (3). In serum, proteolytic processing removes the C-terminal arginine, creating a low activity C5a desArg77 molecule (1). Mouse C5a shares 60% and 82% aa sequence identity to human and rat C5a, respectively. C5a binds to a signaling G-protein coupled receptor (GPCR) (C5aR/CD88), inducing neutrophil chemotaxis and endothelial cell activation (1, 4). It also triggers an oxidative burst in macrophages and neutrophils, and induces release of histamine in basophils and mast cells (1, 4). Alternatively, it may also bind to a non-signaling GPCR termed C5L2 whose function is yet to be determined (5).

## References:

1. Gerard, C. and N.P. Gerard (1994) *Annu. Rev. Immunol.* **12**:775.
2. DiScipio, R.G. *et al.* (1983) *J. Biol. Chem.* **258**:10629.
3. Huber-Lang, M.S. *et al.* (2003) *J. Immunol.* **170**:6115.
4. Gerard, N.P. and C. Gerard (2002) *Curr. Opin. Immunol.* **14**:705.
5. Okinaga, S. *et al.* (2003) *Biochemistry* **42**:9406.