

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-10 in direct ELISAs and Western blots. In Western blots, this antibody does not cross-react with recombinant mouse IL-10.
Source	Monoclonal Mouse IgG _{2B} Clone # 25209
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human IL-10 Ser19-Asn178 Accession # P22301
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS and NaCl with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	5 µg/mL	See Below
Neutralization		Measured by its ability to neutralize IL-10-induced proliferation in the MC/9-2 mouse mast cell line. Thompson-Snipes, L. <i>et al.</i> (1991) <i>J. Exp. Med.</i> 173 :507. The Neutralization Dose (ND ₅₀) is typically 0.005-0.03 µg/mL in the presence of 5 ng/mL Recombinant Human IL-10.

DATA

<p>Western Blot</p> <p>Detection of Human IL-10 by Western Blot. Western blot shows conditioned media of HEK293 human embryonic kidney cell line either mock transfected or transfected with human IL-10. PVDF membrane was probed with 5 µg/mL of Mouse Anti-Human IL-10 Monoclonal Antibody (Catalog # MAB2171) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # HAF018). A specific band was detected for IL-10 at approximately 16 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p>Neutralization</p> <p>Cell Proliferation Induced by IL-10 and Neutralization by Human IL-10 Antibody. Recombinant Human IL-10 (Catalog # 217-L) stimulates proliferation in the MC/9-2 mouse mast cell line in a dose-dependent manner (orange line). Proliferation elicited by Recombinant Human IL-10 (5 ng/mL) is neutralized (green line) by increasing concentrations of Mouse Anti-Human IL-10 Monoclonal Antibody (Catalog # MAB2171). The ND₅₀ is typically 0.005-0.03 µg/mL.</p>
--	---

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

IL-10 is a pleiotropic cytokine that can exert either immunostimulatory or immunosuppressive effects on a variety of cell types. It is produced predominantly by Th2 lymphocytes and other activated lymphoid cells.