

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Histone H2AX when phosphorylated at S139 in Western blots.
<b>Source</b>	Recombinant Monoclonal Rabbit IgG Clone # 2207D
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	Phospho-peptide containing human Histone H2AX S139 site Accession # P16104
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.25 µg/mL	See Below
<b>Intracellular Staining by Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below
<b>CytoF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Histone H2AX is a core histone protein that is phosphorylated at S139 in cells exposed to DNA double-strand break-inducing agents, such as ionizing radiation. The S139 phosphorylated H2AX, termed γ-H2AX, marks the site of DNA double-strand breaks and serves to recruit cell cycle checkpoint and DNA repair factors to the site of damage.