

**DESCRIPTION**

<b>Specificity</b>	Detects TGF-β1 from human, mouse, rat, and other species in direct ELISAs and Western blots. In sandwich ELISAs, less than 2% cross-reactivity with recombinant human (rh) TGF-β3 and recombinant amphibian TGF-β5 and no cross-reactivity with recombinant porcine TGF-β2 or rhTGF-β2 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 9016
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant TGF-β1 and latent TGF-β1
<b>Endotoxin Level</b>	<0.10 EU per 1 µg of the antibody by the LAL method.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


**APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	Recombinant Human TGF-β1 (Catalog # 240-B) under non-reducing conditions only
<b>Immunocytochemistry</b>	3-25 µg/mL	See Below
<b>Immunohistochemistry</b>	8-25 µg/mL	See Below
<b>Intracellular Staining by Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	See Below
<b>Human TGF-β1 Sandwich Immunoassay</b>		<b>Reagent</b>
<b>ELISA Capture</b>	2-8 µg/mL	TGF-β1 Antibody (Catalog # MAB240)
<b>ELISA Detection Standard</b>	0.1-0.4 µg/mL	TGF-β1 Biotinylated Antibody (Catalog # BAF240) Recombinant Human TGF-β1 (Catalog # 240-B)
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	
<b>Neutralization</b>	Measured by its ability to neutralize TGF-β1 inhibition of IL-4-dependent proliferation in the HT-2 mouse T cell line. Tsang, M. <i>et al.</i> (1995) Cytokine 7:389. The Neutralization Dose (ND <sub>50</sub> ) is typically 0.3-1.0 µg/mL in the presence of 0.25 ng/mL Recombinant Human TGF-β1 and 7.5 ng/mL Recombinant Mouse IL-4.	


**DATA**

**Immunocytochemistry**




**TGF-β1 in HEK293 Human Cell Line.** TGF-β1 was detected in immersion fixed HEK293 human embryonic kidney cell line using Mouse Anti-TGF-β1 Monoclonal Antibody (Catalog # MAB240) at 3 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 493-conjugated Anti-Mouse IgG Secondary Antibody (green; Catalog # NL009) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

**Immunohistochemistry**




**TGF-β1 in Human Prostate Cancer Tissue.** TGF-β1 was detected in immersion fixed paraffin-embedded sections of human prostate cancer tissue using Mouse Anti-TGF-β1 Monoclonal Antibody (Catalog # MAB240) at 25 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific labeling was localized to the cytoplasm of epithelial cells in the prostate gland. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

**Intracellular Staining by Flow Cytometry**



**Detection of TGF-β1 in PC-3 Human Cell Line by Flow Cytometry.** PC-3 human prostate cancer cell line was stained with Mouse Anti-TGF-β1 Monoclonal Antibody (Catalog # MAB240, filled histogram) or isotype control antibody (Catalog # MAB002, open histogram), followed by Allophycocyanin-conjugated Anti-Mouse IgG F(ab)<sub>2</sub> Secondary Antibody (Catalog # F0101B). To facilitate intracellular staining, cells were fixed with paraformaldehyde and permeabilized with saponin.

**Neutralization**



**TGF-β1 Inhibition of IL-4-dependent Cell Proliferation and Neutralization by TGF-β1 Antibody.** Recombinant Human TGF-β1 (Catalog # 240-B) inhibits Recombinant Mouse IL-4 (Catalog # 404-ML) induced proliferation in the HT-2 mouse T cell line in a dose-dependent manner (orange line). Inhibition of Recombinant Mouse IL-4 (7.5 ng/mL) activity elicited by Recombinant Human TGF-β1 (0.25 ng/mL) is neutralized (green line) by increasing concentrations of Mouse Anti-TGF-β1 Monoclonal Antibody (Catalog # MAB240). The ND<sub>50</sub> is typically 0.3-1.0 μg/mL.

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

TGF-β1, -2, and -3 are a closely related group of proteins (70-80% sequence homology) that are produced by many cell types and function as growth and differentiation factors. The active forms of TGF-β1, -2, and -3 are disulfide-linked homodimers.