

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human GSK-3 β .
Source	Monoclonal Rat IgG _{2B} Clone # 272535
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human GSK-3 β Met1-Thr420 Accession # P49841
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunocytochemistry	8-25 μ g/mL	See Below

DATA

<p>Immunocytochemistry</p> 	<p>GSK-3β in MCF-7 Human Cell Line. GSK-3β was detected in immersion fixed MCF-7 human breast cancer cell line using 10 μg/mL Human GSK-3β Monoclonal Antibody (Catalog # MAB25061) for 3 hours at room temperature. Cells were stained (red) and counterstained (green). View our protocol for Fluorescent ICC Staining of Cells on Coverslips.</p>
--	---

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Glycogen Synthase Kinase-3 (GSK-3) is a serine/threonine kinase initially identified as an inhibitor of glycogen synthase. Two isoforms (GSK-3 α and GSK-3 β) share 85% amino acid identity. GSK-3 β , inhibited by phosphorylation at S9 by Akt, is involved in energy metabolism, body pattern formation, and neuronal cell development.