

DESCRIPTION


Species Reactivity	Human/Mouse/Rat
Specificity	Detects human MKK3/MKK6 in direct ELISAs and recombinant and endogenous MKK3/MKK6 in Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 275922
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human MKK3 Met1-Ser347 Accession # P46734
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Mitogen-activated protein kinase kinase 3 (MKK3), also known as MEK3, and the closely related MKK6 are activated by proinflammatory cytokines and environmental stresses. Activation occurs through phosphorylation by several upstream MAPK kinase kinases, including ASK1 and TAK1. Both MKK3 and MKK6 are dual specificity protein kinases, phosphorylating and activating the p38 MAP kinases at Thr and Tyr positions within the phosphoacceptor sequence Thr-Gly-Tyr.

PRODUCT SPECIFIC NOTICES

The purchase of this product conveys to the buyer the limited, non-exclusive, non-transferable right (without the right to resell, repackage, or further sublicense) to use these reagents for research purposes only. No other license is granted to the buyer whether expressly, by implication, by estoppel or otherwise. In particular, the purchase of this product does not include nor carry any right or license to use, develop, or otherwise exploit this product commercially, which includes without limitation, provision of services to a third party, generation of commercial databases, clinical diagnostics or therapeutics, or drug development. This product is sold pursuant to a license from the University of Massachusetts, and the University of Massachusetts reserves all other rights to use these reagents. For information on purchasing a license for uses other than research, please contact the University of Massachusetts, Office of Technology Management, 333 South Street, Suite 400, Shrewsbury, Massachusetts, 01545.