

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CXCL5/ENA-78 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) CXCL1, 2, 3, 8, 10, or recombinant mouse CXCL2 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 33160
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human CXCL5/ENA-78 Ala37-Asn114 (predicted) Accession # P42830
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Human CXCL5/ENA-78 (Catalog # 254-XB) under non-reducing conditions only
Immunohistochemistry	8-25 µg/mL	Immersion fixed paraffin-embedded sections of human breast
Neutralization		Measured by its ability to neutralize CXCL5/ENA-78-induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CXCR2. The Neutralization Dose (ND ₅₀) is typically 3-15 µg/mL in the presence of 0.03 µg/mL Recombinant Human CXCL5/ENA-78.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CXCL5, also known as epithelial cell-derived-neutrophil-activating-peptide (ENA78), is an 8 kDa proinflammatory member of the CXC subfamily of chemokines. Its Glu-Leu-Arg (ELR) motif confers angiogenic properties and distinguishes it from ELR⁻ CXC chemokines which are angiostatic. Human CXCL5 shares 57% amino acid (aa) sequence identity with mouse and rat CXCL5. Among other human ELR⁺ chemokines, it shares 77% aa sequence identity with CXCL6/GCP2 and 35%-51% with CXCL1/GRO alpha, CXCL2/GRO beta, CXCL3/GRO gamma, CXCL7/NAP2, and CXCL8/IL8. Inflammatory stimulation upregulates CXCL5 production in multiple hematopoietic cell types, fibroblasts, endothelial cells, and vascular smooth muscle cells. In vivo, CXCL5 is elevated at sites of inflammation and pulmonary fibrosis where it promotes neutrophil infiltration and activation as well as angiogenesis. Its upregulation contributes to increased vascularization, tumor growth, and metastasis in many cancers. Full length CXCL5 (78 aa) is trimmed at the N-terminal end by Cathepsin G and chymotrypsin to ENA74 (74 aa) and ENA70 (70 aa), with the shortened forms showing increased potency relative to full length CXCL5. CXCL5 exerts its effects primarily through interactions with CXCR2. It also binds DARC, a decoy chemokine receptor which can limit CXCR2-mediated responses.