

**DESCRIPTION**

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human IFN- $\gamma$ in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse IFN- $\gamma$ , recombinant rat IFN- $\gamma$ , or recombinant porcine IFN- $\gamma$ is observed. Neutralizes the bioactivity of recombinant human IFN- $\gamma$ and will not neutralize the activity of rat or mouse IFN- $\gamma$ .
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 25723
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human IFN- $\gamma$ Gln24-Gln166 Accession # AAP20098.1
<b>Endotoxin Level</b>	<0.10 EU per 1 $\mu$ g of the antibody by the LAL method.
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 $\mu$ m filtered solution in PBS.


**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Immunocytochemistry</b>	8-25 $\mu$ g/mL	Immersion fixed human peripheral blood mononuclear cells stimulated with PMA and ionomycin
<b>Intracellular Staining by Flow Cytometry</b>	0.25 $\mu$ g/10 <sup>6</sup> cells	See Below
<b>CytoTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	
<b>Neutralization</b>	Measured by its ability to neutralize IFN- $\gamma$ inhibition of EMCV-induced cytopathy in the HeLa human cervical epithelial carcinoma cell line. Meager, A. (1987) in <i>Lymphokines and Interferons, a Practical Approach</i> . Clemens, M. J. <i>et al.</i> (eds): IRL Press. 129. The Neutralization Dose (ND <sub>50</sub> ) is typically 0.2-0.6 $\mu$ g/mL in the presence of 5 ng/mL Recombinant Human IFN- $\gamma$ .	


**DATA**

**Intracellular Staining by Flow Cytometry**



**Detection of IFN- $\gamma$  in Human PBMCs by Flow Cytometry.** Human peripheral blood mononuclear cells (PBMCs) treated with 50 ng/mL PMA, 1  $\mu$ g/mL Ionomycin, and 3  $\mu$ M Monensin overnight were stained with either (A) Mouse Anti-Human IFN- $\gamma$  Monoclonal Antibody (Catalog # MAB2851) or (B) Mouse IgG2B Isotype Control (Catalog # MAB0041) followed by anti-Mouse IgG PE-conjugated secondary antibody (Catalog # F0102B) and Mouse Anti-Human CD3 epsilon APC-conjugated Monoclonal Antibody (Catalog # FAB100A). To facilitate intracellular staining, cells were fixed and permeabilized with FlowX FoxP3 Fixation & Permeabilization Buffer Kit (Catalog # FC012). View our protocol for [Staining Intracellular Molecules](#).

**Neutralization**



**IFN- $\gamma$  Inhibition of EMCV-induced Cytopathy and Neutralization by Human IFN- $\gamma$  Antibody.** Recombinant Human IFN- $\gamma$  (Catalog # 285-IF) reduces the Encephalomyocarditis Virus (EMCV)-induced cytopathy in the HeLa human cervical epithelial carcinoma cell line in a dose-dependent manner (orange line). Inhibition of EMCV activity elicited by Recombinant Human IFN- $\gamma$  (5 ng/mL) is neutralized (green line) by increasing concentrations of Mouse Anti-Human IFN- $\gamma$  Monoclonal Antibody (Catalog # MAB2851). The ND<sub>50</sub> is typically 0.2-0.6  $\mu$ g/mL.

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Interferon-gamma (IFN- $\gamma$ ), also known as type II or immune interferon, exerts a wide range of immunoregulatory activities and is considered to be the prototype proinflammatory cytokine (1, 2). Mature human IFN- $\gamma$  exists as a non-covalently linked homodimer of 20-25 kDa variably glycosylated subunits (3). It shares 90% amino acid (aa) sequence identity with rhesus IFN- $\gamma$ , 59-64% with bovine, canine, equine, feline, and porcine IFN- $\gamma$ , and 37-43% with cotton rat, mouse, and rat IFN- $\gamma$ . IFN- $\gamma$  dimers bind to IFN- $\gamma$  RI (alpha subunits) which then interact with IFN- $\gamma$  RII (beta subunits) to form the functional receptor complex of two  $\alpha$  and two  $\beta$  subunits. Inclusion of IFN- $\gamma$  RII increases the binding affinity for ligand and the efficiency of signal transduction (4, 5). IFN- $\gamma$  is produced by a variety of immune cells under inflammatory conditions, notably by T cells and NK cells (6). It plays a key role in host defense by promoting the development and activation of Th1 cells, chemoattraction and activation of monocytes and macrophages, upregulation of antigen presentation molecules, and immunoglobulin class switching in B cells. It also exhibits antiviral, antiproliferative, and apoptotic effects (6, 7). In addition, IFN- $\gamma$  functions as an anti-inflammatory mediator by promoting the development of regulatory T cells and inhibiting Th17 cell differentiation (8, 9). The pleiotropic effects of IFN- $\gamma$  contribute to the development of multiple aspects of atherosclerosis (7).

**References:**

1. Billiau, A. and P. Matthys (2009) Cytokine Growth Factor Rev. **20**:97.
2. Pestka, S. *et al.* (2004) Immunol. Rev. **202**:8.
3. Gray, P.W. and D.V. Goeddel (1982) Nature **298**:859.
4. Marsters, S.A. *et al.* (1995) Proc. Natl. Acad. Sci. USA **92**:5401.
5. Krause, C.D. *et al.* (2000) J. Biol. Chem. **275**:22995.
6. Schroder, K. *et al.* (2004) J. Leukoc. Biol. **75**:163.
7. McLaren, J.E. and D.P. Ramji (2009) Cytokine Growth Factor Rev. **20**:125.
8. Muhl, H. and J. Pfeilschifter (2003) Int. Immunopharmacol. **3**:1247.
9. Kelchtermans, H. *et al.* (2008) Trends Immunol. **29**:479.