

**DESCRIPTION**

<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects human MIF in direct ELISA and Western Blots.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 932612
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human MIF Met1-Phe114 Accession # P14174
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	2 µg/mL	See Below
<b>Immunohistochemistry</b>	8-25 µg/mL	See Below

**DATA**

<p><b>Western Blot</b></p>	<p><b>Detection of Mouse MIF by Western Blot.</b> Western blot shows lysates of RAW 264.7 mouse monocyte/macrophage cell line and J774A.1 mouse reticulum cell sarcoma macrophage cell line. PVDF membrane was probed with 2 µg/mL of Mouse Anti-Human/Mouse MIF Monoclonal Antibody (Catalog # MAB2892) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # HAF018). A specific band was detected for MIF at approximately 12 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p><b>Immunohistochemistry</b></p> <p><b>MIF in Human Ovary.</b> MIF was detected in immersion fixed paraffin-embedded sections of human ovary using Mouse Anti-Human/Mouse MIF Monoclonal Antibody (Catalog # MAB2892) at 15 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Mouse HRP-DAB Cell &amp; Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific staining was localized to epithelial cells. View our protocol for <a href="#">Chromogenic IHC Staining of Paraffin-embedded Tissue Sections</a>.</p>
----------------------------	---	---

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

MIF (or macrophage migration inhibitory factor) was the first lymphokine/cytokine to be recognized in the pregenomics era (1, 2). Regardless, it is one of the least understood of all inflammatory mediators (1, 3). Human MIF is a 12.5 kDa, 115 amino acid (aa) nonglycosylated polypeptide that is synthesized without a signal sequence (4 - 7). Secretion occurs nonclassically via an ABCA1 transporter (8). The initiating Met is removed, leaving Pro as the first amino acid. The molecule consists of two  $\alpha$ -helices and six  $\beta$ -strands, four of which form a  $\beta$ -sheet. The two remaining  $\beta$ -strands interact with other MIF molecules, creating a trimer (2, 9, 10). Structure-function studies suggest MIF is bifunctional with segregated topology. The N- and C-termini mediate enzyme activity (in theory). Phenylpyruvate tautomerase activity (enol-to-keto) has been demonstrated and is dependent upon Pro at position #1 (11). Amino acids 50 - 65 have also been suggested to contain thiol-protein oxidoreductase activity (12). MIF has proinflammatory cytokine activity centered around aa's 49 - 65. On fibroblasts, MIF induces, IL-1, IL-8 and MMP expression; on macrophages, MIF stimulates NO production and TNF- $\alpha$  release following IFN- $\gamma$  activation (13, 14). MIF apparently acts through CD74 and CD44, likely in some form of trimeric interaction (15, 16). Human MIF is active on mouse cells (14). Human MIF is 90%, 94%, 95%, and 90% aa identical to mouse, bovine, porcine and rat MIF, respectively.

**References:**

1. Norand, E.F. and M. Leech (2005) *Front. Biosci.* **10**:12.
2. Donn, R.P. and D.W. Ray (2004) *J. Endocrinol.* **182**:1.
3. Calandra, T. and T. Roger (2003) *Nat. Rev. Immunol.* **3**:791.
4. Kozak, C.A. *et al.* (1995) *Genomics* **27**:405.
5. Weiser, W.Y. *et al.* (1989) *Proc. Natl. Acad. Sci. USA* **86**:7522.
6. Paralkar, V. and G. Wistow (1994) *Genomics* **19**:48.
7. Wistow, G.J. *et al.* (1993) *Proc. Natl. Acad. Sci. USA* **90**:1272.
8. Flieger, O. *et al.* (2003) *FEBS Lett.* **551**:78.
9. Philo, J.S. *et al.* (2004) *Biophys. Chem.* **108**:77.
10. Sun, H-W. *et al.* (1996) *Protein Eng.* **9**:631.
11. Stamps, S.L. *et al.* (2000) *Biochemistry* **39**:9671.
12. Nguyen, M.T. *et al.* (2003) *J. Biol. Chem.* **278**:33654.
13. Sato, A. *et al.* (2003) *Dev. Comp. Immunol.* **27**:401.
14. Bernhagen, J. *et al.* (1994) *Biochemistry* **33**:14144.
15. Leng, L. *et al.* (2003) *J. Exp. Med.* **197**:1467.
16. Meyer-Siegler, K.L. and P.L. Vera (2005) *J. Urol.* **173**:615.