

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Granzyme A in direct ELISAs.
Source	Recombinant Monoclonal Mouse IgG ₁ Clone # 356428R
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Granzyme A Cys26-Val262 Accession # P12544
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

ELISA This antibody functions as an ELISA capture antibody when paired with Goat Anti-Human Granzyme A Antigen Affinity-purified Polyclonal Antibody (Catalog # [AF2905](#)).

This product is intended for assay development on various assay platforms requiring antibody pairs. We recommend the Human Granzyme A DuoSet ELISA Kit (Catalog # [DY2905-05](#)) for convenient development of a sandwich ELISA.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Granzyme A is a member of the granzyme family of the serine proteases found specifically in the cytotoxic granules of cytotoxic T lymphocytes (CTL) and natural killer (NK) cells. Granzyme A is the most abundant protease in CTL and NK cells. It induces caspase-independent cell death when introduced into target cells by perforin (1). Human granzyme A is synthesized as a 262 amino acid (aa) precursor with a signal peptide (aa 1-26), a propeptide (aa 27-28) and a mature chain (aa 29-262) (2). After being activated by lysyl endopeptidase removal of the propeptide, Granzyme A cleaves thioester substrates. Mature human granzyme A shares 71% amino acid sequence identity with mouse granzyme A.

References:

1. Lieberman, J. and Z. Fan (2003) *Curr. Opin. Immunol.* **15**:553.
2. Gershenfeld, H.K. *et al.* (1988) *Proc. Natl. Acad. Sci. USA* **85**:1184.