

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human MICL/CLEC12A in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) CLEC1, 2, 2A, 3B, 9A, 10A, 12B, 14A, rhCD302/CLEC13A recombinant mouse (rm) CLEC4B2, or rmMICL is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 687317
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human MICL/CLEC12A Thr67-Ala265 Accession # Q5QGZ9
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Human MICL (myeloid inhibitory C-type lectin-like receptor; also CLL-1 and KLRL-1) is a 75 kDa member of the group V C-type lectin-like receptor family of molecules. Group V molecules have a lectin-type domain that binds non-sugar ligands. MICL is a 265 amino acid (aa) type II transmembrane (TM) glycoprotein that contains a 200 aa extracellular domain (ECD). Multiple isoforms of MICL are reported. One potentially utilizes an alternate start site that adds 10 aa to the cytoplasmic domain. There is one potential soluble form that lacks the TM segment, and two truncated ECD isoforms. Human MICL ECD is 53%, 65%, and 55% aa identical to the ECD in mouse, dog and cow, respectively.