

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CREB in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 321120
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human CREB Lys136-Asp341 Accession # P16220
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	8-25 µg/mL	See Below

DATA

Immunohistochemistry



CREB in Human Prostate. CREB was detected in immersion fixed paraffin-embedded sections of human spinal prostate array using Human CREB Monoclonal Antibody (Catalog # MAB2989) at 15 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

The cAMP response element binding protein (CREB), also known as CREB1, belongs to the bZIP family of transcription factors. Each bZIP family member contains a basic domain that mediates DNA binding and a leucine zipper domain that facilitates dimerization. Within the promoter of target genes, CREB dimers bind cAMP response elements, defined by the palindromic consensus sequence TGACGTCA.