

## DESCRIPTION


<b>Species Reactivity</b>	Human/Mouse/Rat
<b>Specificity</b>	Detects endogenous human, mouse and rat PP1 $\alpha$ . Based on the amino acid homologies, the antibody is expected to cross-react with PP1 $\beta$ and PP1 $\gamma$ .
<b>Source</b>	Monoclonal Mouse IgG <sub>2A</sub> Clone # 319319
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human PP1 $\alpha$ catalytic subunit Lys41-Pro298 Accession # P62136
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 $\mu$ m filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 $\mu$ g/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Protein Phosphatase 1, also known as PP1 and PPP1, is an enzyme that removes phosphate groups attached to serine or threonine residues in proteins. The holoenzyme is composed of two subunits, a catalytic subunit (PP1 $\alpha$ ) that is highly conserved throughout evolution, and a wide variety of regulatory subunits that target the enzyme to specific subcellular compartments and proteins. The catalytic subunit is ubiquitously expressed in tissues and cell lines.