

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human COMP in Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 2127-F5
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Bacteria-derived recombinant human COMP
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	2 µg/mL	See Below
Immunoprecipitation	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human COMP/Thrombospondin-5 (Catalog # 3134-CP), see our available Western blot detection antibodies

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Cartilage Oligomeric Matrix Protein (COMP), also known as Thrombospondin-5, is a 110 kDa multidomain calcium binding protein that associates with other extracellular matrix molecules. Thrombospondin-1 and -2 constitute subgroup A and form homotrimers, whereas Thrombospondin-3, -4, and COMP constitute subgroup B and form homopentamers (1-4). The human COMP cDNA encodes a 757 amino acid (aa) precursor that includes a 20 aa signal sequence followed by a non-collagenous coiled-coil domain, four EGF-like repeats, seven TSP type-3 repeats, and a globular TSP C-terminal domain (5). Human COMP shares 86-93% aa sequence identity with rat, mouse, equine, bovine, and canine COMP. Within the TSP type-3 repeats and TSP C-terminal domain, human COMP shares 60%, 61%, 74%, and 80% aa sequence identity with human Thrombospondin-1, -2, -3, and -4, respectively. The coiled coil domain mediates the association of COMP into disulfide-linked homopentamers with a central hub and peripheral globular domains connected by flexible strands (6, 7). An axial pore is formed by the coiled coil assembly and binds vitamin D₃ which is involved in bone and cartilage metabolism (8). An RGD sequence in the third TSP type-3 repeat mediates chondrocyte attachment *via* Integrin α5β1, although when reduced and in the absence of calcium, attachment is mediated *via* Integrin αVβ3 (9). COMP is upregulated in rheumatoid arthritis and osteoarthritis, hepatocellular carcinomas, chronic pancreatitis, and pancreatic carcinomas (10-12). Elevated circulating COMP levels are used as a biomarker for early onset of some skeletal disorders (10). Several mutations are associated with skeletal dysplasias, and the most common, a point mutation in the third TSP type-3 repeat, results in diminished calcium binding ability (13, 14).

References:

1. Adams, J.C. and J. Lawler (2004) *Int J. Biochem. Cell Biol.* **36**:961.
2. Posey, K.L. *et al.* (2004) *Int. J. Biochem. Cell Biol.* **36**:1005.
3. Adams, J.C. (2004) *Int. J. Biochem. Cell Biol.* **36**:1102.
4. Mann, H.H. *et al.* (2004) *J. Biol. Chem.* **279**:25294.
5. Newton, G. *et al.* (1994) *Genomics*, **24**:435.
6. DiCesare, P. *et al.* (1995) *J. Orthopaedic Res.* **13**:422.
7. Efimov, V.P. *et al.* (1994) *FEBS Lett.* **341**:54.
8. Ozbek, S. *et al.* (2002) *EMBO J.* **21**:5960.
9. Chen, F.H. *et al.* (2005) *J. Biol. Chem.* **280**:32655.
10. Wislowska, M. and B. Jablonska (2005) *Clin. Rheumatol.* **24**:278.
11. Xiao, Y. *et al.* (2004) *J. Gastroenterol. Hepatol.* **19**:296.
12. Liao, Q. *et al.* (2003) *Scand. J. Gastroenterol.* **38**:207.
13. Kennedy, J. *et al.* (2005) *Eur. J. Hum. Genet.* **13**:547.
14. Hou, J. *et al.* (2000) *Cell Calcium* **27**:309.