

Human Wnt-9a Antibody


Monoclonal Mouse IgG₁ Clone # 326711 Catalog Number: MAB3157

| DESCRIPTION | | | |
|--------------------|---|--|--|
| Species Reactivity | Human | | |
| Specificity | Detects human Wnt-9a in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human Wnt-2, -3a, -6, -7 -7b, recombinant mouse Wnt-1, -2b, -4, -5a, -5b, -8a, -8b, -9b, -10a, -10b, -11, or -16 is observed. | | |
| Source | Monoclonal Mouse IgG ₁ Clone # 326711 | | |
| Purification | Protein A or G purified from hybridoma culture supernatant | | |
| Immunogen | E. coli-derived recombinant human Wnt-9a Pro48-Glu92 & Arg231-Phe304 Accession # O14904.2 | | |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS. | | |

APPLICATIONS


Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

| | Recommended Concentration | Sample | |
|--|--|-----------|--|
| Western Blot | 5 μg/mL | See Below | |
| Intracellular Staining by Flow Cytometry | 2.5 µg/10 ⁶ cells | See Below | |
| CyTOF-ready | Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation. | | |



Detection of Human Wnt-9a by Western Blot.

Western blot shows lysates of KATO-III human gastric carcinoma cell line and ME-180 human cervical epithelial carcinoma cell line. PVDF membrane was probed with 5 µg/mL Mouse Anti-Human Wnt-9a Monoclonal Antibody (Catalog # MAB3157) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # HAF007). For additional reference, recombinant human Wnt-9a (20 ng) was included. Specific bands for native Wnt-9a were detected at approximately 30 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 6.



Detection of Wnt-9a in MCF-7 Human Cell Line by Flow Cytometry. MCF-7 human breast cancer cell line was stained with Mouse Anti-Human Wnt-9a Monoclonal Antibody (Catalog # MAB3157, filled histogram) or isotype control AA175antibody (Catalog # MAB002, open histogram), followed by Allophycocyanin-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # F0101B). To facilitate intracellular staining, cells were fixed with paraformaldehyde and permeabilized with saponin.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 0.5 mg/mL in sterile PBS.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Wnt-9a (Wingless related integration site family member 9a, formerly Wnt-14) is a member of the Wnt family, which are ligands for frizzled family receptors. Wnt-9a is a secreted glycoprotein expressed in skeletal muscle and heart as well as in B-CLL and gastric, pancreatic and breast cancer cells. Human Wnt-9a shares 98% aa sequence identity with mouse and rat Wnt-9a.

Rev. 2/7/2018 Page 1 of 1

