



DESCRIPTION	
<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human CTRP5/C1qTNF5 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) CTRP1 or rhCTRP4 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 332923
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human CTRP5/C1qTNF5 Ser16-Ala243 (Gln44Arg) Accession # Q9BXJ0
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**  
Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 µg/mL	Recombinant Human CTRP5/C1qTNF5
<b>Immunohistochemistry</b>	5-25 µg/mL	See Below
<b>Intracellular Staining by Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA	
<p><b>Immunohistochemistry</b></p>  <p><b>CTR5/C1qTNF5 in Human Liver.</b> CTR5/C1qTNF5 was detected in immersion fixed paraffin-embedded sections of human liver using Mouse Anti-Human CTR5/C1qTNF5 Monoclonal Antibody (Catalog # MAB3167) at 10 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Mouse HRP-DAB Cell &amp; Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific staining was localized to sinusoids. View our protocol for <a href="#">Chromogenic IHC Staining of Paraffin-embedded Tissue Sections</a>.</p>	<p><b>Intracellular Staining by Flow Cytometry</b></p>  <p><b>Detection of CTR5/C1qTNF5 in A172 Human Cell Line by Flow Cytometry.</b> A172 human glioblastoma cell line was stained with Mouse Anti-Human CTR5/C1qTNF5 Monoclonal Antibody (Catalog # MAB3167, filled histogram) or isotype control antibody (Catalog # MAB0041, open histogram), followed by Allophycocyanin-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # F0101B). To facilitate intracellular staining, cells were fixed with Flow Cytometry Fixation Buffer (Catalog # FC004) and permeabilized with Flow Cytometry Permeabilization/Wash Buffer I (Catalog # FC005). View our protocol for <a href="#">Staining Intracellular Molecules</a>.</p>

PREPARATION AND STORAGE	
<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**  
CTR5, also known as C1qTNF5, belongs to the highly conserved family of Acrp30/Adiponectin paralogs known as C1q and TNF-related protein family. All family members share a modular organization comprising an N-terminal signal peptide, a short variable region, a collagenous domain and a C-terminal globular domain. CTRP proteins are predicted to have trimeric structures that assemble into hexameric and higher order molecular forms. Human and mouse CTRP5 share 94% amino acid sequence identity.