

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human OX40/TNFRSF4 in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant human (rh) 4-1BB, rhBAFF R, rhCD27, rhCD30, rhCD40, rhDR3, rhDR6, rhEDAR, rhFAS, rhGITR, rhHVEM, rhLTβR, rhNGF R, rhRANK, rhRELT, rhTAJ, rhTNF RI, rhTNF RII, rhTRAIL, rhTRAIL R4, rhTRAIL R3, rhTWEAK R, or rhXEDAR is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 443318
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human OX40/TNFRSF4 Leu29-Ala216 Accession # P43489
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 μg/mL	Recombinant Human OX40/TNFRSF4 Fc Chimera (Catalog # 3388-OX)
<b>Flow Cytometry</b>	0.25 μg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

OX40 (CD134; TNFRSF4) is a T cell co-stimulatory molecule of the TNF receptor superfamily that coordinates with other membrane-bound co-stimulators such as CD28, CD40, CD30, CD27 and 4-1BB (1-3). OX40 is expressed on naïve CD4<sup>+</sup> T cells only after engagement of the TCR by antigen presenting cells (APC; dendritic and B cells), and co-stimulation by CD40/CD40 ligand and CD28/B7. It is maximal at 2-5 days post activation, or 4 hours post reactivation of memory T cells (3-6). Human OX40 is a 48 kDa type I transmembrane glycoprotein with a 28 amino acid (aa) signal sequence, a 185 aa extracellular domain (ECD) that has four TNFR-Cys repeats and an O-glycosylated hinge region, a 20 aa transmembrane segment, and a 41 aa cytoplasmic domain (3). The ECD of human OX40 shows 71%, 68%, 67%, 64% and 64% aa identity with feline, canine, rabbit, mouse and rat OX40 ECD, respectively. Engagement of OX40 on activated CD4<sup>+</sup> T cells by OX40 ligand on activated dendritic cells promotes T cell survival and proliferation, prolongs the immune response, and enhances the number of cells making the transition from effector to memory T cells (1-6). OX40 signal transduction includes binding TNF receptor-associated factors (TRAFs), and activating NFκB and PI3 kinase to enhance expression of cytokines, antiapoptotic Bcl-2 family members, survivin and the chemokine receptor CXCR5 (5-8). CXCR5 promotes T cell migration to germinal centers to deliver B cell help (5). Studies using knockout or transgenic mice, and agonistic or blocking antibodies, show that OX40/OX40L interaction is critical for establishing or reactivating memory T cells and breaking immune tolerance (9). Blockade of OX40 engagement is efficacious in animal models of allergic airway inflammation, graft-versus-host disease and autoimmune disease (10-14).

**References:**

1. Salek-Ardakani, S. and M. Croft (2006) *Vaccine* **24**:872.
2. Hori, T. (2006) *Int. J. Hematol.* **83**:17.
3. Latza, U. *et al.* (1994) *Eur. J. Immunol.* **24**:677.
4. Murata, K. *et al.* (2000) *J. Exp. Med.* **191**:365.
5. Fillatreau, S and D. Gray (2003) *J. Exp. Med.* **197**:195.
6. Gramaglia, I. *et al.* (1998) *J. Immunol.* **161**:6510.
7. Rogers, P.R. *et al.* (2001) *Immunity* **15**:445.
8. Song, J. *et al.* (2005) *Immunity* **22**:621.
9. Bansal-Pakala, P. *et al.* (2001) *Nat. Med.* **7**:907.
10. Salek-Ardakani, S. *et al.* (2003) *J. Exp. Med.* **198**:315.
11. Jember, A. G. *et al.* (2001) *J. Exp. Med.* **193**:387.
12. Demirci, G. *et al.* (2004) *J. Immunol.* **172**:1691.
13. Blazar, B.R. *et al.* (2003) *Blood* **101**:3741.
14. Higgins, L.M. *et al.* (1999) *J. Immunol.* **162**:486.