

Human/Mouse/Rat MKP-3 Antibody

Monoclonal Mouse IgG_{2A} Clone # 363246 Catalog Number: MAB3576

DESCRIPTION		
Species Reactivity	Human/Mouse/Rat	
Specificity	Detects endogenous human, mouse, and rat MKP-3 in Western blots. In Western blots, this antibody does not cross-react with recombinant human MKP-1 or MKP-2.	
Source	Monoclonal Mouse IgG _{2A} Clone # 363246	
Purification	Protein A or G purified from hybridoma culture supernatant	
Immunogen	E. coli-derived recombinant human MKP-3 Ile2-Thr381 Accession # Q16828	
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.	


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	1 μg/mL	See Below

DATA

Western Blot



Detection of Human and Mouse MKP-3 by Western Blot. Western Blot shows lysates of HeLa human cervical epithelial carcinoma cell line, C2C12 mouse myoblast cell line, and CTLL-2 mouse cytotoxic T cell line. PVDF membrane was probed with 1 µg/mL of Human/Mouse/Rat MKP-3 Monoclonal Antibody (Catalog # MAB3576) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # HAF007). A specific band was detected for MKP-3 at approximately 42 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
	*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- e Ose a manual derrost freezer and avoid repeated freeze-tha
 - 12 months from date of receipt, -20 to -70 °C as supplied.
 1 month, 2 to 8 °C under sterile conditions after reconstitution.
 - 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

MAP Kinase Phosphatase 3 (MKP-3), also called PYST1 and Dual Specificity Phosphatase 6 (DUSP6), dephosphorylates serine, threonine, and tyrosine residues in proteins. On Western blots, MKP-3 has an apparent molecular weight of 42 kilodaltons. It has a high degree of substrate selectivity for dephosphorylating ERK kinases over stress-related kinases such as p38 and JNK. MKP-3 is ubiquitinylated prior to degradation in the proteosome. The rate of MKP-3 degradation is enhanced by ERK-mediated phosphorylation.

Rev. 2/7/2018 Page 1 of 1

