

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CCL19/MIP-3 β in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 948918
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human CCL19/MIP-3 β Gly22-Ser98 Accession # Q99731
Endotoxin Level	<0.10 EU per 1 μ g of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Neutralization Measured by its ability to neutralize CCL19/MIP-3 β -induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CCR7. The Neutralization Dose (ND50) is typically 0.1-0.5 μ g/mL in the presence of 50 ng/mL Recombinant Human CCL19/MIP-3 β .

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

MIP-3 β , also known as ELC (EBI1-Ligand Chemokine), is one of many novel β chemokines identified through bioinformatics. MIP-3 β cDNA encodes a 98 amino acid (aa) residue precursor protein with a predicted 21 aa residue signal peptide that is cleaved to form the 77 aa residue mature secreted protein. MIP-3 β is distantly related to other β chemokines (20-30% aa sequence identity) and the gene for MIP-3 β has been mapped to chromosome 9p13 rather than chromosome 17 where the genes for many human β chemokines are clustered. MIP-3 β has been shown to be constitutively expressed in various lymphoid tissues (including thymus, lymph nodes, appendix and spleen). The expression of MIP-3 β is down-regulated by the anti-inflammatory cytokine IL-10. MIP-3 β has been shown to be a unique functional ligand for CCR7 (previously referred to as the Epstein-Barr virus-induced gene 1 (EBI1) orphan receptor), a chemokine receptor that is expressed in various lymphoid tissues and activated B and T lymphocytes. EBI1 is strongly up-regulated in B cells infected with Epstein-Barr virus and T cells infected with herpesvirus 6 or 7.