


DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human CCL15/MIP-1δ in ELISAs and Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 52130
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human CCL15/MIP-1δ Gln22-Ile113 Accession # Q16663.2
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Human CCL15/MIP-1δ 92 aa (Catalog # 363-MG)
Human CCL15/MIP-1δ Sandwich Immunoassay		Reagent
ELISA Capture	2-8 µg/mL	Human CCL15/MIP-1δ Antibody (Catalog # MAB363)
ELISA Detection	0.1-0.4 µg/mL	Human CCL15/MIP-1δ Biotinylated Antibody (Catalog # BAF363)
Standard		Recombinant Human CCL15/MIP-1δ 92 aa (Catalog # 363-MG)
Neutralization		Measured by its ability to neutralize CCL15/MIP-1δ-induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CCR1. The Neutralization Dose (ND ₅₀) is typically 3-15 µg/mL in the presence of 16 ng/mL Recombinant Human CCL15/MIP-1δ 92 aa.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CCL15, a human CC chemokine, was isolated from a human fetal spleen cDNA library. CCL15 cDNA encodes a predicted 113 amino acid (aa) protein containing a putative signal peptide of 21 aa that is cleaved to generate a 92 aa residue mature protein. Within the CC family members, human CCL15 shares 45%, 44%, 35%, and 30% aa homology with mouse C10, human MPIF-1, human HCC-1, and mouse MIP-1 γ , respectively. The gene for MIP-1 δ is found on chromosome 17 where the genes for most of the human CC chemokines are located. Human CCL15 is expressed in T and B lymphocytes, NK cells, monocytes and monocyte-derived dendritic cells. Human MIP-1 δ is chemotactic for T cells and monocytes and has been shown to induce calcium flux in human CCR-1-transfected cells.

References:

1. Wang, W. *et al.* (1998) J. Clinical Immunol. **18**:214