

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Complement Component C1rLP in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) C1r, rhC1s, rhHaptoglobin, or rhMASP3 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 393608
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Complement Component C1rLP Ser41-Asn487 Accession # AA156781
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Human Complement Component C1rLP
Immunoprecipitation	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human Complement Component C1rLP, see our available Western blot detection antibodies

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Complement Component 1, r subcomponent-like protein (C1rLP) is a serine protease that is mainly expressed in the liver and has structural similarity to C1r, a component of the classical complement activation pathway. C1rLP activates pro-C1s and pro-haptoglobin. The amino acid sequence of human C1rLP is 99% and 75% identical to that of chimpanzee and dog/mouse/rat, respectively.