

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Lck in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 355713
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Lck
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS and NaCl with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below

DATA

Immunocytochemistry



Lck in Human PBMCs. Lck was detected in immersion fixed human peripheral blood mononuclear cells (PBMCs) using Mouse Anti-Human Lck Monoclonal Antibody (Catalog # MAB3704) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). View our protocol for [Fluorescent ICC Staining of Non-adherent Cells](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Lck (p56lck) is a 56 kDa cytosolic phosphoprotein within the Src family of non-receptor tyrosine kinases. The 509 amino acid (aa) human Lck contains an SH3 domain (aa 61-121), an SH2 domain (aa 127-224) and a protein kinase domain (aa 251-259). Within aa 2-66, human Lck shares 89% aa sequence identity with mouse and rat Lck. A short (363 aa) isoform contains alternate sequence starting at aa 348, while a 539 aa isoform contains inserted sequence after aa 321. Lck interacts with T cell CD4 and CD8 molecules and plays a pivotal role in regulating T cell activation. It can be overexpressed in cancers and functions as an oncogene.