

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects human and mouse MafK in Western blots. In Western blots, no cross-reactivity with recombinant human (rh) MafB or rhMafF is observed. In Western blots, prolonged film exposures may slightly detect rhMafG.
Source	Monoclonal Mouse IgG _{2B} Clone # 381923
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human MafK Thr2-Ser156 Accession # O60675
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	2 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Maf family members form a unique subclass of basic-leucine zipper (bZIP) transcription factors. Maf proteins are subdivided into two groupings: large, including c-Maf, Nrl, MafA, and MafB; and small, including MafF, MafG, and MafK, also known as p18. Large Mafs contain an N-terminal acidic domain important for transcriptional activation that is lacking in small Maf family members. Heterodimers of MafK and p45 NF-E2, another bZIP factor, promote globin gene expression and erythroid differentiation in erythroleukemia cells.