

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects endogenous human and mouse Hck at 59 kDa in Western blots.
Source	Monoclonal Rat IgG _{2A} Clone # 394903
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Hck Met22-Asp80 Accession # P08631
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Hemopoietic cell kinase (Hck) is a member of the Src family of tyrosine kinases. Hck expression is prominent in cells of myeloid lineage, particularly in granulocytes and monocytes, with protein levels increasing during terminal differentiation and macrophage activation. Hck contains a unique amino-terminal domain that is subject to lipid modifications, targeting this kinase to the plasma membrane.