

DESCRIPTION


| | |
|---------------------------|---|
| Species Reactivity | Human |
| Specificity | Detects human FCAR/CD89 in direct ELISAs. In direct ELISAs, no cross-reactivity with Fcγ RIA, RIIA, or RIIB is observed. |
| Source | Monoclonal Mouse IgG ₁ Clone # 488032 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | NS0 mouse myeloma cell line transfected with human FCAR/CD89 Gln22-Lys287 Accession # P24071 |
| Formulation | Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μm filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|--|---------------|
| Flow Cytometry | 0.25 μg/10 ⁶ cells | See Below |
| CyTOF-ready | Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation. | |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Reconstitution | Reconstitute at 0.5 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

FCAR, also called FcαRI or CD89, is a variably glycosylated 50-100 kDa myeloid-specific type I transmembrane (TM) Fc receptor for IgA that is a member of the multichain immune recognition receptor (MIRR) family (1-3). Human FCAR contains a 21 amino acid (aa) signal sequence and extracellular (ECD), TM and cytoplasmic domains of 206, 19 and 41 aa, respectively (4). Arg230 within the TM domain supports interaction with the ITAM-containing signaling subunit, FcRγ, which contains a TM Asp (5-7). Two ECD C2-type Ig-like domains (EC1 and 2) are oriented at right angles (8). Up to two molecules of FCAR can bind one molecule of serum IgA via EC1 (8). Many splice variants have been reported, but only two have been identified as proteins (9, 10). The α.2 form, which lacks 22 aa just prior to the TM domain, is exclusively expressed in alveolar macrophages. The α.3 form lacks EC2. FCAR binds monomeric, polymeric and secretory IgA, but does not mediate the barrier function of secretory IgA in mucosal epithelium (1-3). Shedding and circulation of polymeric IgA/FCAR immune complexes has been reported (11). Circulating neutrophils, eosinophils, and monocytes express FCAR (12). Tissue expression of FCAR is mainly from neutrophils; FCAR is downregulated as monocytes differentiate to tissue macrophages (12). On neutrophils, a significant amount of FCAR lacks FcRγ, but can still be endocytosed to early endosomes and recycled to the cell surface (5). Binding of serum IgA to FCAR is transient and anti-inflammatory, inhibiting IgG or IgE-induced degranulation (6). Sustained aggregation of FCAR results in inflammatory responses (6). FcRγ signaling is required for these and for transport to late endosomes (5-7). Human FCAR shows 55-58% aa identity with rat, horse and cow FCAR. No ortholog occurs in mouse. FCAR structure resembles the KIR/ILT/LIR/MIR family more than other IgA receptors, including plgR, Fcα/μR, asialoglycoprotein receptor (ASGR1) and transferrin receptor (TfR) (1-3).

References:

1. Wines, B. D. and P. M. Hogarth (2006) *Tissue Antigens* **68**:103.
2. Otten, M. A. and M. van Egmon (2004) *Immunol. Lett.* **92**:23.
3. Montiero, R. C. and J. G. J. van de Winkel (2003) *Annu. Rev. Immunol.* **21**:177.
4. Maliszewski, C. R. *et al.* (1990) *J. Exp. Med.* **172**:1665.
5. Launay, P. *et al.* (1999) *J. Biol. Chem.* **274**:7216.
6. Pasquier, B. *et al.* (2005) *Immunity* **22**:31.
7. Shen, L. *et al.* (2001) *Blood* **97**:205.
8. Herr, Y. *et al.* (2003) *Nature* **423**:614.
9. Patry, C. *et al.* (1996) *J. Immunol.* **156**:4442.
10. Togo, S. *et al.* (2003) *FEBS Lett.* **535**:205.
11. van der Boog, P. J. M. *et al.* (2002) *J. Immunol.* **168**:1252.
12. Hamre, R. *et al.* (2003) *Scand. J. Immunol.* **57**:506.