

DESCRIPTION


Species Reactivity	Human
Specificity	Detects endogenous human Bcl-9 in Western blots.
Source	Monoclonal Mouse IgG _{2B} Clone # 417514
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Bcl-9 Met1009-Gly1328 Accession # O00512
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Bcl-9 (B-cell lymphoma 9; also Protein legless homolog) is a 149 kDa transcriptional regulator that belongs to the Bcl-9 family of proteins. It is expressed in multiple tissues and serves to recruit Pygopus to the Wnt-pathway β-catenin-TCF complex in the nucleus. Bcl-9 and Bcl-9-2 are considered evolutionary duplicates of Legless that perform the same task with different regulatory mechanisms. Human Bcl-9 is 1426 amino acids (aa) in length. It contains one phosphothreonine and three phosphoserine sites, two poly-Pro regions (aa 514-517 and 970-973), and one poly-Ala segment (aa 900-903). There is one potential alternate start site at Met27, and a variant isoform exists that shows a four aa substitution for aa 1391-1426. Over aa 1009-1328, human Bcl-9 is 96% aa identical to mouse Bcl-9. B cell cancers often have translocations at the 3'UTR region of the Bcl-9 gene.