

#### DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse IL-9 in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant human IL-9 is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 222604
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant mouse IL-9 Gln19-Pro144 Accession # P15247.1
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	Recombinant Mouse IL-9 (Catalog # 409-ML)
<b>Intracellular Staining by Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	Mouse splenocytes treated with anti-CD3/anti-CD28, Recombinant Mouse IL-4 (Catalog # 404-ML), Human TGF-β1 (Catalog # 100-B), and PMA/Ionomycin to induce Th9 development.
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

#### PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

#### BACKGROUND

Mouse IL-9 was originally identified as a T cell-derived T cell growth factor III/P40 which could support the long term growth of certain mouse T helper clones in the absence of antigen or antigen-presenting cells. IL-9 can also prolong the *in vitro* survival of other T cell clones as well as potentiate the IL-2 dependent proliferation of mouse fetal thymocytes. However, this cytokine has no growth-stimulating activity on mouse cytolytic T cell clones or fresh T cells. Mouse IL-9 also has mast cell enhancing activity (MEA) and can enhance the mIL-3- or mIL-4-dependent proliferation of mouse bone marrow-derived mast cells. Furthermore, IL-9 will synergize with erythropoietin to support erythroid colony formation *in vitro*. The mouse IL-9 cDNA encodes a 144 amino acid residue precursor protein with an 18 amino acid signal peptide that is cleaved to form the mature cysteine-rich protein with a predicted molecular mass of 14 kDa. Mouse IL-9 contains four potential N-linked glycosylation sites and the native mIL-9 is a highly glycosylated protein. Human and mouse IL-9 share 56% amino acid sequence homology. Although mouse IL-9 is active on human cells, human IL-9 is not active on mouse cells.

#### References:

1. Renauld, J.E. *et al.* (1995) *J. Leukoc. Biol* **57**:303.