

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human S100A13 in ELISAs and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant mouse S100A13 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 854114
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human S100A13 Ala2-Lys98 Accession # Q99584
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	2 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

S100A13 is an 11 kDa member of the S100 (soluble in 100% saturated ammonium sulfate) family of vertebrate EF-hand Ca<sup>++</sup>-binding proteins (1-3). It is widely expressed as a homodimer with two 98 amino acid (aa) long subunits (2, 3). Human S100A13 shares 83%, 90%, 91%, 87%, 78% and 47% aa identity with mouse, rat, cow, dog, opossum and chicken S100A13, respectively. Like other S100 proteins, S100A13 is small and generally acidic, but contains a basic residue-rich sequence at the C terminus, and two EF hand motifs that bind with Ca<sup>++</sup> differing affinities (2-4). Some S100 proteins, including S100A13, are able to bind the cell surface receptor for advanced glycation end-products (RAGE) (5). Despite lacking a signal sequence, S100A13 plays an important role in Cu<sup>++</sup>-dependent export of FGF-1 (FGF acidic) and IL-1 $\alpha$  from the cell in response to stresses such as heat shock, anoxia and starvation (6-8). Binding of copper is necessary for formation of a multi-protein complex between S100A13, FGF-1 and p40 synaptotagmin-1 (syt-1) (9, 10). Cu<sup>++</sup> ions supplied by S100A13 are thought to oxidize and downregulate the activity of FGF-1 prior to export (10). Calcium influx may also play a similar role in FGF-1 release from neuronal cells (11). S100A13 is composed of four amphiphilic helices that may interact with acidic phospholipid headgroups. With FGF-1 and syt-1, S100A13 likely perturbs the membrane, which allows the S100A13 protein complex to exit the cell (4, 12). S100A13 has been proposed as a marker for angiogenesis in tumors and endometrium, due to its role in stress-induced export of FGF-1 (13, 14). Based on in house studies, S100A13 has also been found to promote neurite outgrowth from rat cortical embryonic neurons (15).

## References:

1. Santamaria-Kisiel, L. *et al.* (2006) *Biochem. J.* **396**:201.
2. Wicki, R. *et al.* (1996) *Biochem. Biophys. Res. Commun.* **227**:594.
3. Ridinger, K. *et al.* (2000) *J. Biol. Chem.* **275**:8686.
4. Li, M. *et al.* (2007) *Biochem. Biophys. Res. Commun.* **356**:616.
5. Hsieh, H.-L. *et al.* (2004) *Biochem. Biophys. Res. Commun.* **316**:949.
6. Landriscina, M. *et al.* (2001) *J. Biol. Chem.* **276**:22544.
7. Sivaraja, V. *et al.* (2006) *Biophys. J.* **91**:1832.
8. Mandinova, A. *et al.* (2003) *J. Cell Sci.* **116**:2687.
9. Prudovsky, I. *et al.* (2002) *J. Cell Biol.* **158**:201.
10. Landriscina, M. *et al.* (2001) *J. Biol. Chem.* **276**:25549.
11. Matsunaga, H. and H. Ueda (2006) *Cell. Mol. Neurobiol.* **26**:237.
12. Graziani, I. *et al.* (2006) *Biochem. Biophys. Res. Commun.* **349**:192.
13. Landriscina, M. *et al.* (2006) *J. Neurooncol.* **80**:251.
14. Hayrabyan, S. *et al.* (2005) *Reprod. Biol.* **5**:51.
15. R&D Systems (2007) In-house data.