

DESCRIPTION


Species Reactivity	Human/Mouse/Rat
Specificity	Detects endogenous human, mouse, and rat PON3 in Western blots. In Western blots, this antibody does not cross-react with recombinant human (rh) PON1 or rhPON2.
Source	Monoclonal Rat IgG _{2A} Clone # 454929
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human PON3 Ala30-Leu354 Accession # AAH70374
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

The paraoxonase (PON) gene family of antioxidant enzymes includes three known members located adjacent to each other on chromosome 7. Paraoxonase/lactonase 3 (PON3) is a 354 amino acid, 39.6 kDa protein that is secreted into the bloodstream and associates with high-density lipoprotein (HDL). PON3 has lactonase activity (EC 3.1.1) and rapidly hydrolyzes lactones. PON3 exhibits no paraoxonase and very limited arylesterase activities. PON3 is an important antioxidant and can inhibit the oxidation of low-density lipoprotein (LDL), a function that may slow the initiation and progression of atherosclerosis. PON3 is glycosylated at Asn323 and alternatively spliced variants encoding different protein isoforms have been described; however, only one has been fully characterized. Human PON3 shares 81% amino acid identity with mouse and rat PON3.