

**DESCRIPTION**



<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse CD151. Stains mouse CD151 transfectants but not irrelevant transfectants.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 455807
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	HEK293 human embryonic kidney cell line transfected with mouse CD151 Met1-Tyr253 Accession # BAA22447
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below
<b>Immunocytochemistry</b>	8-25 µg/mL	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**

<p><b>Flow Cytometry</b></p>  <p><b>Detection of CD151 in bEnd.3 Mouse Cell Line by Flow Cytometry.</b> bEnd.3 mouse endothelioma cell line was stained with Rat Anti-Mouse CD151 Monoclonal Antibody (Catalog # MAB4609, filled histogram) or isotype control antibody (Catalog # MAB006, open histogram), followed by Phycoerythrin-conjugated Anti-Rat IgG F(ab)<sub>2</sub> Secondary Antibody (Catalog # F0105B).</p>	<p><b>Immunocytochemistry</b></p>  <p><b>CD151 in bEnd.3 Mouse Cell Line.</b> CD151 was detected in immersion fixed bEnd.3 mouse endothelioma cell line using Rat Anti-Mouse CD151 Monoclonal Antibody (Catalog # MAB4609) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. View our protocol for <a href="#">Fluorescent ICC Staining of Cells on Coverslips</a>.</p>
--	---

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

CD151 is a 29 kDa palmitoylated glycoprotein in the tetraspanin superfamily and is expressed by endothelial cells, epithelial cells, megakaryocytes, and platelets. It associates with other tetraspanin proteins and with some integrins including α3/β1, α6/β1, α6/β4, and α7/β1. CD151 modulates cellular adhesion, cell migration, and platelet activation. Mouse CD151 shares 94%-96% aa sequence identity with human and rat CD151.