

DESCRIPTION



Species Reactivity	Human
Specificity	Stains human GPR34 transfectants but not irrelevant transfectants.
Source	Monoclonal Mouse IgG _{2A} Clone # 419859
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with human GPR34 Accession # Q9UPC5
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below
Intracellular Staining by Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA

<p>Immunocytochemistry</p>  <p>GPR34 in A172 Human Cell Line. GPR34 was detected in immersion fixed A172 human glioblastoma cell line using Mouse Anti-Human GPR34 Monoclonal Antibody (Catalog # MAB4617) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). View our protocol for Fluorescent ICC Staining of Cells on Coverslips.</p>	<p>Intracellular Staining by Flow Cytometry</p>  <p>Detection of GPR34 in A172 Human Cell Line by Flow Cytometry. A172 human glioblastoma cell line was stained with Mouse Anti-Human GPR34 Monoclonal Antibody (Catalog # MAB4617, filled histogram) or isotype control antibody (Catalog # MAB003, open histogram), followed by Phycoerythrin-conjugated Anti-Mouse IgG F(ab)₂ Secondary Antibody (Catalog # F0102B). To facilitate intracellular staining, cells were fixed with paraformaldehyde and permeabilized with saponin.</p>
--	--

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

GPR34 is a widely expressed 75-90 kDa seven transmembrane segment glycoprotein. It is upregulated on microglia during inflammation and functions as a receptor for lysophosphatidyl-L-serine on mast cells to promote degranulation. Multiple isoforms of GPR34 may result from multiple translation initiation sites and alternative splicing. Human GPR34 shares 89% amino acid sequence identity with mouse and rat GPR34.