

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human NAALADase-like 2/NAALADL2 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 817225
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human NAALADase-like 2/NAALADL2 Ser152-Asn795 Accession # Q58DX5
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

NAALADL2 (N-acetylated α-linked acidic dipeptidase like 2) is a member of the peptidase M28 family of enzymes. It is presumably a type II transmembrane (TM) protein that may have O-glycosyl hydrolase activity. Human NAALADL2 is 795 amino acids (aa) in length. It is believed to contain a cytoplasmic and TM segment at the N-terminus, followed by a peptidase domain (aa 444-596) and a TIR-like dimerization region (aa 688-777). There are multiple splice forms. One shows a 4 aa substitution for aa 292-795, a second shows an alternate start site at Met18 with a 24 aa substitution for aa 314-795, and a third contains an alternate start site at Met283, accompanied by a deletion of aa 363-411 and 633-795. Over aa 152-795, human NAALADL2 shares 87% and 82% aa sequence identity with mouse and canine NAALADL2, respectively.