

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human Integrin $\beta 8$ in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human Integrins $\beta 1$, $\beta 2$, $\beta 3$, $\beta 4$, $\beta 5$, $\beta 6$, $\beta 7$, or recombinant mouse Integrin $\beta 8$ is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 416922
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Integrin $\beta 8$ Glu43-Arg684 (predicted) Accession # P26012
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25 μ g/10 ⁶ cells	See Below
CytoF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Integrin beta 8 (Integrin $\beta 8$) is a 90 kDa type I transmembrane glycoprotein of the Integrin family of adhesion molecules. It associates with Integrin αV to form a receptor for vitronectin, fibrin, and the latency associated peptide (LAP). Binding to LAP promotes the proteolytic release of active TGF- β from LAP. Integrin $\alpha V\beta 8$ is required for vascular morphogenesis in the embryonic brain and yolk sac. Within the extracellular domain, human Integrin $\beta 8$ shares 87% aa sequence identity with mouse and rat Integrin $\beta 8$.