

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human 5T4 in direct ELISAs. In direct ELISAs, 100% cross-reactivity with recombinant mouse 5T4 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 524731
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human 5T4 Ser31-Ser355 Accession # Q13641
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	8-25 µg/mL	See Below

DATA

Immunohistochemistry



5T4 in Human Placenta. 5T4 was detected in immersion fixed paraffin-embedded sections of human placenta using Mouse Anti-Human 5T4 Monoclonal Antibody (Catalog # MAB4975) at 15 µg/mL overnight at 4 °C. Before incubation with the primary antibody tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific labeling was localized to syncytiotrophoblast cells. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Human 5T4 (oncofetal antigen 5T4; also TPBG and trophoblast glycoprotein) is a 72 kDa glycoprotein member of the LRR family of proteins. It is expressed on trophoblasts, tumor cells, ovarian cuboidal epithelium and embryonic stem cells, and impacts cell adhesion and motility. The human 5T4 cDNA encodes a type I transmembrane protein precursor that is 420 amino acids (aa) in length. It contains a 324 aa extracellular region (aa 32-355) that shows one Ser-rich region followed by seven Leu-rich repeats (aa 90-355). Over aa 31-355, human 5T4 shares 81% and 85% aa identity with mouse and canine 5T4, respectively.