

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human LRRC4 in direct ELISAs. In direct ELISAs, greater than 25% cross-reactivity with recombinant human (rh) NGL-1/LRRC4C is observed, and no cross-reactivity with rhLRRC4B is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 701419
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human LRRC4 Ala39-Lys527 Accession # Q9HBW1
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	8-25 µg/mL	See Below

DATA

Immunohistochemistry



LRRC4 in Human Brain. LRRC4 was detected in immersion fixed paraffin-embedded sections of human brain (medulla) using Mouse Anti-Human LRRC4 Monoclonal Antibody (Catalog # MAB49951) at 15 µg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific staining was localized to neuronal cell bodies, processes, and synaptic puncta. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

LRRC4 (Leucine rich repeat/LRR-containing protein 4), also called NGL-2 (netrin-G ligand-2) or NAG14 (nasopharyngeal carcinoma-associated gene 14) is a 55 kDa (predicted, unglycosylated) type I transmembrane protein that is a member of the NGL family of synaptic LRR adhesion molecules (1, 2). Human LRRC4 cDNA encodes 653 amino acids (aa) that include a 38 aa signal sequence, a 489 aa extracellular domain (ECD), a 21 aa transmembrane domain, and a 105 aa cytoplasmic domain. The ECD contains nine LRRs (aa 74-288), a C2 type Ig like domain (aa 354-440), and a Thr-rich segment (aa 455-526). Within the ECD, human LRRC4 shares 98% aa identity with mouse and rat, 99% aa identity with canine and bovine, and 99.6% aa identity with equine LRRC4. It also shares 54-55% aa identity with family members LRRC4C/NGL-1 and LRRC4B/NGL-3, but each recognizes different ligands (1). LRRC4 is predominantly expressed in the brain on neurons and astrocytes as a ligand for netrin-G2 on the dendritic surface of synaptic neurons (2-4). It is proposed to regulate the formation of excitatory synapses via recruitment of PSD-95 to the cytoplasmic domain after aggregation of LRRC4 at the surface (3, 5). It suppresses proliferation by downregulating cell signaling pathways, resulting in altered expression of cell cycle regulating proteins and delay at the late G1 phase (1, 2, 6-8). It is thus considered a tumor suppressor protein and is often downregulated in brain tumors, particularly gliomas (1, 2, 6). Forced expression of LRRC4 in tumor cells slows proliferation and promotes differentiation (1, 4, 9). Addition of soluble LRRC4 to cultured neurons reduces excitatory synapse formation (3).

References:

1. Woo, J. *et al.* (2009) *Mol. Cell. Neurosci.* **42**:1.
2. Zhang, Q. *et al.* (2005) *FEBS Lett.* **579**:3674.
3. Kim, S. *et al.* (2006) *Nat. Neurosci.* **9**:1294.
4. Wu, M. *et al.* (2007) *Acta Biochim Biophys Sin (Shanghai)* **39**:731.
5. Nishimura-Akiyoshi, S. *et al.* (2007) *Proc. Natl. Acad. Sci. USA* **104**:14801.
6. Wu, M. *et al.* (2006) *Mol. Biol. Cell* **17**:3534.
7. Wu, M. *et al.* (2008) *J. Cell. Biochem.* **103**:245.
8. Wu, M. *et al.* (2008) *J. Cell. Physiol.* **214**:65.
9. Zhang, W. *et al.* (2008) *Genes Brain Behav.* **7**:385.