



DESCRIPTION	
<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse BMP-4 in direct ELISAs and Western blots.
<b>Source</b>	Recombinant Monoclonal Rabbit IgG Clone # 1128D
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived mature mouse BMP-4 Ser293-Arg408 Accession # P21275
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS		
<b>Please Note:</b> Optimal dilutions should be determined by each laboratory for each application. <i>General Protocols</i> are available in the <i>Technical Information</i> section on our website.		
	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.25 µg/mL	See Below
<b>Immunohistochemistry</b>	1-15 µg/mL	See Below

DATA	
<p><b>Western Blot</b></p>  <p><b>Detection of Mouse BMP-4 by Western Blot.</b> Western blot shows Recombinant Mouse BMP-4 (Catalog # 5020-BP) and lysates of mouse embryo tissue. PVDF membrane was probed with 0.25 µg/mL of Rabbit Anti-Mouse BMP-4 Monoclonal Antibody (Catalog # MAB5020) followed by HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog # HAF008). Specific bands were detected for BMP-4 at approximately 40-55 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p><b>Immunohistochemistry</b></p>  <p><b>BMP-4 in Mouse Embryo.</b> BMP-4 was detected in immersion fixed frozen sections of mouse embryo (15 d.p.c.) using Rabbit Anti-Mouse BMP-4 Monoclonal Antibody (Catalog # MAB5020) at 2 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Rabbit HRP-DAB Cell &amp; Tissue Staining Kit (brown; Catalog # CTS005) and counterstained with hematoxylin (blue). Specific staining was localized to cytoplasm in gastrointestinal tract cells. View our protocol for <a href="#">Chromogenic IHC Staining of Frozen Tissue Sections</a>.</p>

PREPARATION AND STORAGE	
<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

BMP-4 is a TGF- $\beta$  superfamily ligand that is widely expressed from early embryogenesis through adulthood. It plays an important role in mesenchyme formation, epidermal determination, suppression of neural induction, the development of multiple organs, and tissue repair (1-5). The mouse BMP-4 precursor contains a 273 amino acid (aa) propeptide and a 116 aa mature protein (6). The propeptide is cleaved intracellularly by furin or proprotein convertase 6, enabling the 15 kDa mature BMP-4 monomer to form an active disulfide linked homodimer or heterodimer with BMP-7 (7-9). Mature mouse and human BMP-4 share 98% aa sequence identity. Mouse BMP-4 shares 85% aa sequence identity with mouse BMP-2 and 35%-54% with other mouse BMPs. Compared to BMP-4 homodimers, BMP-4/BMP-7 heterodimers exhibit a greater potency in inducing osteogenic differentiation (9). In *Xenopus*, the heterodimers can also induce the formation of mesoderm, whereas BMP-4 homodimers only provide ventralizing signals for existing mesoderm (10). BMP-4 signals through tetrameric complexes composed of type I (primarily Activin RIA or BMPR-IA) and type II (primarily Activin RIIA or BMPR-II) receptors (11, 12). The bioavailability of BMP-4 is regulated by its interaction with multiple proteins and glycosaminoglycans (13-15).

**References:**

1. Zhang, P. *et al.* (2008) *Blood* **111**:1933.
2. Gambaro, K. *et al.* (2006) *Cell Death Differ.* **13**:1075.
3. Simic, P. and S. Vukicevic (2005) *Cytokine Growth Factor Rev.* **16**:299.
4. Sadlon, T.J. *et al.* (2004) *Stem Cells* **22**:457.
5. Frank, D.B. *et al.* (2005) *Circ. Res.* **97**:496.
6. Oida, S. *et al.* (1995) *DNA Seq.* **5**:273.
7. Cui, Y. *et al.* (1998) *EMBO J.* **17**:4735.
8. Cui, Y. *et al.* (2001) *Genes Dev.* **15**:2797.
9. Aono, A. *et al.* (1995) *Biochem. Biophys. Res. Commun.* **210**:670.
10. Nishimatsu, S. and G.H. Thomsen (1998) *Mech. Dev.* **74**:75.
11. Chen, D. *et al.* (2004) *Growth Factors* **22**:233.
12. Lavery, K. *et al.* (2008) *J. Biol. Chem.* April 24 epub.
13. Rosen, V. (2006) *Ann. N.Y. Acad. Sci.* **1068**:19.
14. Jones, C.M. and J.C. Smith (1998) *Dev. Biol.* **194**:12.
15. Takada, T. *et al.* (2003) *J. Biol. Chem.* **278**:43229.