

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CCL23/Ckβ8-1 in ELISAs and Western blots. In sandwich immunoassays, no cross-reactivity with human CCL23/Ckβ8 isoform is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 103952
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human CCL23/Ckβ8-1 Met46-Asn137 Accession # AAD10846
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Human CCL23/Ckβ8-1 aa 46-137 (Catalog # 508-CK)
Human CCL23/Ckβ8-1 Sandwich Immunoassay		Reagent
ELISA Capture	2-8 µg/mL	Human CCL23/Ckβ8-1 Antibody (Catalog # MAB508)
ELISA Detection	0.1-0.4 µg/mL	Human CCL23/Ckβ8-1 Biotinylated Antibody (Catalog # BAF508)
Standard		Recombinant Human CCL23/Ckβ8-1 aa 46-137 (Catalog # 508-CK)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CCL23 (SCY A23), a splice variant of CKβ8 (also known as MIPF-1), is a β chemokine isolated from the THP-1 cDNA library. CCL23 cDNA encodes a 137 amino acid (aa) residue precursor protein with a putative 21 aa residue signal peptide. Compared to CKβ8-1, CKβ8/MIPF-1 lacks a 17 aa residue stretch (Leu47-Gly63) present in CCL23 and has a unique arginine at residue 46. CKβ8-1/CKβ8 (MIPF-1) and leukotactin/MIP-1δ form a subgroup of β chemokines that have 6 conserved cysteine residues and an extended amino-terminus preceding the conserved cysteine pair. CCL23 shares approximately 73% nucleotide sequence identity with leukotactin (MIP-1δ), and the two genes have been localized to human chromosome 17 within 200 kb of each other. CCL23 mRNA expression has been detected in the pancreas, heart and skeletal muscle. Both CKβ8-1 and CKβ8 (MIPF-1) are potent agonists of CCR1 and have been shown to chemoattract peripheral blood lymphocytes and monocytes. The two chemokines have also been reported to chemoattract neutrophils and to inhibit colony formation by human hematopoietic progenitor cells. Similar to CKβ8 (MIPF-1), the truncated form of CKβ8-1 (aa 46-132) produced at R&D Systems has been shown to have greatly enhanced biological activity compared to the full length mature CKβ8-1 (aa 22-132).

References:

1. Youn, B-S. *et al.* (1998) *Blood* **91**:3118.
2. Macphee, C.H. *et al.* (1998) *J. Immunol.* **161**:6273.